


Mitochondrial Permeability Uncouples Elevated Autophagy and Lifespan Extension

Graphical Abstract



Authors

Ben Zhou, Johannes Kreuzer, Caroline Kumsta, ..., Malene Hansen, Wilhelm Haas, Alexander A. Soukas

Correspondence

asoukas@mgh.harvard.edu

In Brief

The role of autophagy in lifespan extension depends on modulation of mitochondrial permeability via the action of the kinase SGK1.

Highlights

- SGK1 regulates autophagy in both *C. elegans* and mammalian cells
- Elevated autophagy and mPTP opening shorten lifespan in *sgk-1*/mTORC2 mutant worms
- SGK-1 phosphorylates mPTP component VDAC1 on Ser104, promoting its degradation
- Loss of SGK function exaggerates mPTP-dependent hepatic ischemia/reperfusion injury

Mitochondrial Permeability Uncouples Elevated Autophagy and Lifespan Extension

Ben Zhou,^{1,2,3} Johannes Kreuzer,⁴ Caroline Kumsta,⁵ Lianfeng Wu,^{1,2,3} Kimberli J. Kamer,^{3,6} Lucydalila Cedillo,^{1,2,3} Yuyao Zhang,^{1,2,3} Sainan Li,^{1,2,3} Michael C. Kacergis,^{1,2,3} Christopher M. Webster,^{1,2,3} Geza Fejes-Toth,⁷ Aniko Naray-Fejes-Toth,⁷ Sudeshna Das,⁸ Malene Hansen,⁵ Wilhelm Haas,⁴ and Alexander A. Soukas^{1,2,3,9,*}

¹Department of Medicine, Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA 02114, USA

²Department of Medicine, Harvard Medical School, Boston, MA 02115, USA

³Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA

⁴Center for Cancer Research, Massachusetts General Hospital, Boston, MA 02114, USA

⁵Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA 92037, USA

⁶Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA

⁷Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth, Hanover, NH 03755, USA

⁸MGH Biomedical Informatics Core and Department of Neurology, Massachusetts General Hospital, Boston, MA 02114, USA

⁹Lead Contact

*Correspondence: asoukas@mgh.harvard.edu

<https://doi.org/10.1016/j.cell.2019.02.013>

SUMMARY

Autophagy is required in diverse paradigms of lifespan extension, leading to the prevailing notion that autophagy is beneficial for longevity. However, why autophagy is harmful in certain contexts remains unexplained. Here, we show that mitochondrial permeability defines the impact of autophagy on aging. Elevated autophagy unexpectedly shortens lifespan in *C. elegans* lacking serum/glucocorticoid regulated kinase-1 (*sgk-1*) because of increased mitochondrial permeability. In *sgk-1* mutants, reducing levels of autophagy or mitochondrial permeability transition pore (mPTP) opening restores normal lifespan. Remarkably, low mitochondrial permeability is required across all paradigms examined of autophagy-dependent lifespan extension. Genetically induced mPTP opening blocks autophagy-dependent lifespan extension resulting from caloric restriction or loss of germline stem cells. Mitochondrial permeability similarly transforms autophagy into a destructive force in mammals, as liver-specific *Sgk* knockout mice demonstrate marked enhancement of hepatocyte autophagy, mPTP opening, and death with ischemia/reperfusion injury. Targeting mitochondrial permeability may maximize benefits of autophagy in aging.

INTRODUCTION

A hallmark of aging is the accumulation of molecular damage, dysfunctional organelles, and defective enzymes (López-Otín et al., 2013). Autophagy is an ancient stress response present in all living eukaryotic cells, responsible for clearing intracellular damaged proteins, protein aggregates, and dysfunctional

organelles. Autophagy is generally thought of as a positive longevity modulatory factor. An unbiased screen for chronological aging factors in yeast identified multiple short-lived mutants with defects in autophagy (Matecic et al., 2010). In organisms such as the nematode *C. elegans*, autophagy is also critical for increased longevity evident in insulin-signaling mutants, during caloric restriction, and loss of germline stem cells (Hansen et al., 2018). In mice, disruption of the Beclin 1-BCL2 complex increases autophagy and promotes longevity (Fernández et al., 2018).

In spite of these generally beneficial effects, autophagy may also play a detrimental role in health, promoting diseases such as cancer, diabetes, neurodegeneration, and ischemic injury (Thorburn, 2014). With extended starvation, high levels of autophagy promote death in *C. elegans* (Kang et al., 2007). These observations imply that cellular factors and context determine whether autophagy is beneficial or detrimental to health.

Mitochondrial oxidative stress and high cellular calcium concentrations trigger opening of the mitochondrial permeability transition pore (mPTP). Opening of the mPTP, which is regulated by the voltage-dependent anion channel (VDAC), the adenine nucleotide translocators (Ant proteins), BAX, PiC, F₁F₀ ATP synthase, and cyclophilin D, permits diffusion of molecules of <1,500 Da to flow freely across the mitochondrial inner membrane (Kwong and Molkentin, 2015). Transient mPTP opening regulates calcium and mitochondrial reactive oxygen species (ROS) signaling, while prolonged mPTP opening results in mitochondrial energetic dysfunction, organelle swelling, rupture, and apoptotic or necrotic cell death (Kwong and Molkentin, 2015). Recent evidence suggests that opening of the pore also plays a role in mitochondrial quality control, as mPTP opening depolarizes mitochondria, marking them for autophagic clearance (Gottlieb and Mentzer, 2010). However, direct and mechanistic connections of the mPTP with regulation of autophagy have not been made.

The heteromeric protein kinase mTOR complex 2 (mTORC2) controls cellular metabolism, proliferation, and survival by phosphorylating members of the AGC (PKA/PKG/PKC) family




Figure 1. mTORC2 Regulates Autophagy in Both *C. elegans* and Mouse Hepatocytes

(A) SGK-1::GFP detected by fluorescence microscopy. Scale bar, 200 μ m (n = 67 worms for control and 69 for starvation, Student's t test).
 (B) *C. elegans* *sgk-1* mutants show increased autolysosomes (arrows) by electron microscopy versus WT. Scale bar, 500 nm (n = 59 and 87 for WT and *sgk-1*, respectively, Student's t test).
 (C) Increased autophagy levels in seam cells of *rict-1* and *sgk-1* mutants by GFP::LGG-1 puncta (n = 70, 73, and 71 for WT, *rict-1*, and *sgk-1*, respectively, one-way ANOVA).
 (D) Western blotting of LGG-1::GFP, LGG-1::GFP-PE, and cleaved free GFP protein levels in fed and starved *rict-1* and *sgk-1* mutants (n = 8 biological replicates, one-way ANOVA).
 (E) Fluorescence microscopy of GFP::LGG-1 puncta in the proximal intestinal cells of *rict-1* and *sgk-1* mutants expressing SGK1 or lipidation defective mutant LGG-1(G116A).
 (F and G) Quantification of GFP::LGG-1 puncta in the intestine, muscle, hypodermis, and pharynx of *rict-1* (F) and *sgk-1* (G) mutants expressing LGG-1 or LGG-1(G116A) (n > 25 cells analyzed per group, two-way ANOVA).
 (H) Western blotting of LGG-1::GFP and free GFP protein levels in AML12 hepatocytes treated with vehicle (veh) or chloroquine (CQ).
 (I and J) Western blotting of SGK1 and LC3 protein levels in AML12 hepatocytes treated with HBSS or CQ.

(legend continued on next page)

of protein kinases, including serum/glucocorticoid regulated kinase 1 (SGK1) (García-Martínez and Alessi, 2008). SGK1 is also regulated at different levels by growth factors, glucocorticoids, and inflammatory factors and has been implicated in responses to oxidative stress, apoptosis, and DNA damage (Di Cristofano, 2017). In worms, *sgk-1* is a major effector of metabolic regulation downstream of mTORC2. *ric1-1*, the *C. elegans* ortholog of the essential mTORC2 component Rictor, and *sgk-1* modulate lifespan in a complex manner, depending on food source and temperature (Mizunuma et al., 2014; Soukas et al., 2009). Molecular determinants of longevity, metabolism, and development acting downstream of mTORC2 and *sgk-1* remain incompletely understood.

Here, we show that in spite of shortened lifespan in *ric1-1* and *sgk-1* mutants under standard laboratory conditions, these animals have increased levels of autophagy. Reducing function of autophagy genes restores normal lifespan in mTORC2 mutants, indicating that elevated autophagy is detrimental rather than beneficial in the context of dysfunctional mTORC2 signaling to SGK-1. We establish that the reason that autophagy shortens lifespan is due to a concomitant increase in mitochondrial permeability in mTORC2 mutants, attributable to increased VDAC1 protein level. We find that even in wild-type (WT) *C. elegans*, increased mitochondrial permeability shortens lifespan. Most importantly, this mechanism generalizes to many long-lived genetic mutants dependent upon autophagy. Under each circumstance tested, induction of mitochondrial permeability completely reverses autophagy-dependent lifespan extension. Finally, we show that hepatic ischemia/reperfusion (I/R) injury is enhanced in liver-specific *Sgk* knockout mice due to increased mPTP opening. Thus, enhanced autophagy in the setting of increased mitochondrial permeability is a detrimental combination conserved from nematodes to mammals.

RESULTS

mTORC2 Pathway Members RICT-1 and SGK-1 Negatively Regulate Autophagy

Levels of a SGK-1::GFP reporter decrease in the starved state, suggesting that reduced SGK-1 signaling participates in the metabolic adaptation to starvation (Figures 1A and S1A). Phosphorylation of α catalytic subunit AMP-activated protein kinase Thr172, which is elevated in starvation, is increased in *sgk-1* mutants (Figure S1B). *sgk-1* mutant worms deliver a reduced brood size over an extended period (Figure S1C), similar to animals that are calorically restricted (Crawford et al., 2007; Wang et al., 2014). These results suggest that a key role of SGK-1 is to suppress adaptive responses to starvation in the fed state.

We hypothesized that mTORC2 and SGK-1 may function as negative regulators of catabolic processes such as autophagy during nutrient sufficiency. In line with this, the number of autophagic vesicles evident by electron microscopy and GFP::LGG-1 puncta (*C. elegans* ATG8 ortholog, which marks autophagosomes) is increased in fed *ric1-1* and *sgk-1* mutant worms (Figures 1B and 1C). Western blotting confirms increased LGG-1 protein levels and LGG-1 processing in *ric1-1* and *sgk-1* mutant worms under both fed and starved conditions (Figure 1D). Numbers of GFP::LGG-1 puncta are increased in intestine, muscle, hypodermis, and pharynx in *ric1-1* mutant worms; in *sgk-1* mutant worms, numbers of GFP::LGG-1 puncta are increased in intestine and hypodermis but not in muscle and pharynx (Figures 1E–1G). This suggests that *sgk-1* modulates intestinal and hypodermal autophagy downstream of *ric1-1*. We confirmed that GFP::LGG-1-positive puncta represent autophagosomes and not unspecific aggregates by using mutant GFP-tagged LGG-1(G116A) protein that is expected to be defective in lipidation and autophagosome-membrane targeting (Manil-Ségalen et al., 2014). In GFP::LGG-1(G116A) reporter animals, no increased puncta were seen in *ric1-1* mutant and *sgk-1* mutant worms (Figures 1E–1G).

Increased numbers of GFP::LGG-1 puncta can be seen either in the setting of increased autophagic flux or, alternatively, due to the inhibition of late stages of autophagy (Zhang et al., 2015). We investigated autophagic flux with chloroquine (CQ) treatment, which increases lysosomal pH and impairs lysosomal function (Klionsky et al., 2016). CQ treatment induces accumulation of GFP::LGG-1 and free GFP protein in *ric1-1* and *sgk-1* mutants versus WT worms (Figure 1H). This indicates increased autophagic flux in *ric1-1* and *sgk-1* mutants and increased GFP::LGG-1 puncta seen in these animals are not due to a block in late steps of autophagy.

To explore whether *Sgk* also modulates autophagy in mammals, we generated two distinct loss of function models for *Sgk1*: CRISPR/Cas9 *Sgk1* knockout AML12 hepatocytes and liver-specific *Sgk1* knockout mice (hereafter referred to as *Sgk1*^{Lko}) (Fejes-Tóth et al., 2008). Autophagy initiation is increased in both *Sgk1* knockout AML12 hepatocytes and primary *Sgk1*^{Lko} hepatocytes, as indicated by both immunoblotting for LC3A/B conversion (LC3A/B I to LC3A/B II) (Figures 1I and S1E) and CYTO-ID staining autophagic vacuoles (Figures S1D and S1F). Finally, CQ treatment in *Sgk1* knockout AML12 hepatocytes leads to LC3A/B II accumulation, indicating increased autophagic flux (Figure 1J).

Inhibition of Autophagy Restores Normal Lifespan in Short-Lived mTORC2 Mutants

Given the strong positive association of elevated autophagy and longevity, we evaluated the role of autophagy in *C. elegans*

(H) GFP::LGG-1 and free GFP levels in WT, *ric1-1*, and *sgk-1* mutants under vehicle and chloroquine (CQ, 100 mM for 18 h) treatment (n = 5 biological replicates, two-way ANOVA).

(I) Autophagy in *Sgk1* knockout (*Sgk1*($-/-$)) AML12 hepatocytes as indicated by LC3A/B turnover in both full medium and Hank's balanced salt solution (HBSS) starvation conditions (n = 6 for control and n = 5 HBSS, two-way ANOVA).

(J) LC3A/B conversion in *Sgk1*($-/-$) versus WT AML12 cells with or without 50 μ M chloroquine (CQ) for 4 h (n = 10 for vehicle and n = 6 for CQ, two-way ANOVA).

*p < 0.05, **p < 0.01. All bars indicate means and SEM.

See also Figure S1.




Figure 2. Inhibition of Autophagy Restores Normal Lifespan in Short-Lived *sgk-1* and *rict-1* Mutants

(A–D) RNAi of *bec-1* in (A) *sgk-1* and (C) *rict-1* mutants and RNAi of *lgg-1* in (B) *sgk-1* and (D) *rict-1* mutants restores normal lifespan versus wild-type (WT) worms treated with empty vector RNAi (ev) (log rank test).

(E) GFP::LGG-1 puncta in the intestine are reduced by *bec-1* RNAi in *sgk-1* mutants. Scale bar, 20 μ m (n = 45 worms for ev and n = 46 for *bec-1*, Student's t test).

(F and G) RNAi of *unc-51* partially restores normal lifespan in *sgk-1* (F) and *rict-1* (G) mutants (log rank test).

(H) HLH-30::GFP protein level in *sgk-1* mutants and WT animals. Scale bar, 200 μ m (n = 20 worms for WT and n = 15 for *sgk-1*, Student's t test).

(I) Two different RNAi targeting *hlh-30* both extend lifespan in *sgk-1* and *rict-1* mutants (log rank test).

For tabular survival data and biological replicates see also Table S1. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. All bars indicate means and SEM.

See also Figure S2.

mTORC2 and *sgk-1* mutant lifespan. Surprisingly, RNAi of autophagy components *bec-1* (*C. elegans* *Beclin1* homolog) and *lgg-1* restored normal lifespan in short-lived *sgk-1* and *rict-1* mutants (Figures 2A–2D). Confirming that *bec-1* RNAi inhibits autophagy, GFP::LGG-1 puncta decrease in *sgk-1* mutants after *bec-1* RNAi (Figure 2E). RNAi of *bec-1* extends lifespan similarly in both *sgk-1* mutants and *rict-1*;*sgk-1* double mutants (Figure S2A), suggesting *sgk-1* is the main modulator of lifespan downstream of *rict-1*. RNAi of the autophagy regulator *unc-51* (*C. elegans* *ULK1* homolog) also partially restored lifespan and

nearly completely normalized the low brood size of *sgk-1* mutant worms (Figures 2F, 2G, and S2B). Silencing of autophagy components also accelerates *sgk-1* mutant growth (Figure S2C). In contrast, RNAi of these components in WT worms had no effect on development and modestly shortens lifespan (Figures 2A–2D and S2D). RNAi of autophagy genes has no effect on fat levels in *sgk-1* mutants (Figure S2E).

C. elegans HLH-30, homolog of mammalian transcription factor EB, is a positive transcriptional regulator of autophagy (Lapierre et al., 2013). Levels of HLH-30 are elevated in *sgk-1* mutants

as indicated by a HLH-30::GFP reporter (Figure 2H). As with *bec-1* RNAi, knockdown of *hlh-30* significantly extends the lifespan of *sgk-1* mutants (Figure 2I).

Because these results are opposite to the positive effect of enhanced autophagy in many long-lived mutants, we surmised that the cellular context in mTORC2 mutants renders autophagy detrimental rather than beneficial. To begin to elucidate this context, we evaluated the spatiotemporal requirements for *bec-1*. Consistent with our prior observation that mTORC2 regulates metabolism, growth, and reproduction via action in *C. elegans* intestine (Soukas et al., 2009), intestine-specific *bec-1* RNAi restores normal lifespan and slow development in *rict-1* mutant worms (Figures S2F–S2H). Whole life *bec-1* knockdown had the greatest quantitative effect on restoration of normal lifespan in *sgk-1* mutants, versus larval and adult only knockdowns (Figures S2I and S2J and RNAi efficiency in S2K). Thus, intestinal autophagy in all life stages is detrimental to *sgk-1* mutant longevity.

SGK-1 Regulates mPTP Opening

We employed unbiased protein mass spectrometry in order to determine the mechanistic tie between SGK-1 and autophagy. We used GFP-tagged SGK-1 transgenic *C. elegans* in order to co-immunoprecipitate (coIP) SGK-1 interacting proteins, versus a GFP-only transgenic strain. Analysis of proteins that coIP with SGK-1 indicated nearly the entire complement of protein regulators of the mPTP: VDAC-1, ANT-1.1, ANT-1.3, SPG-7, F-ATP synthase components ATP-5 and F58F12.1, and phosphate carrier protein SLC25A3 homolog F01G4.6 (Kwong and Molkentin, 2015; Shanmughapriya et al., 2015). The mPTP is a critical regulator of mitochondrial homeostasis, and it initiates autophagy under serum starvation or glucagon treatment (Elmore et al., 2001). To confirm that the mPTP and SGK-1 interact, we selected as bona fide SGK interactors only proteins that (1) have two or more unique peptide matches, and (2) are not present in GFP control or have at least 5-fold the number of total peptides and 5-fold the average precursor intensity versus GFP control. Using these criteria, six mPTP proteins interact with SGK-1. These six mPTP proteins are overrepresented in the total set of 244 SGK interactors (from our mass spectrometry data), assuming 1,150 mitochondrial proteins as the universe (mitoCarta 2.0) (Calvo et al., 2016) (hypergeometric test p value = 0.000113).

We evaluated direct protein-protein interaction between SGK-1 and regulators of the mPTP. GST-pull down suggests that the direct interaction of SGK-1 with the mPTP occurs through binding with VDAC-1 (Figure 3A). Interaction between mammalian SGK1 and VDAC1 was confirmed by endogenous coIP (Figures 3B and 3C) and reciprocal coIP of overexpressed SGK1 and VDAC1 (Figure S3A) in 293T cells and is not affected by PI3K pathway inhibition (Figure S3B). Overall, these data are in line with our and others' findings that SGK1 localizes to mitochondria (Cordas et al., 2007).

Because SGK1 interacts with VDAC, we next asked whether SGK1 modulates mitochondrial permeability by regulating mPTP opening. As assessed with the calcium retention capacity assay, the mPTP opens after significantly fewer calcium pulses in AML12 hepatocytes lacking *Sgk1* (Figures 3D and S3C). These data suggest that a normal role of SGK1 is to maintain mitochon-

drial homeostasis by supporting mPTP closure and mitochondrial calcium buffering capacity.

We next evaluated whether aberrant opening of the mPTP in *sgk-1* mutant *C. elegans* is deleterious to longevity. Indeed, RNAi of the mPTP regulating gene *ant-1.1* in *rict-1* and *sgk-1* mutants completely reverses their shortened lifespan and moderately extends lifespan of WT worms (Figure 3E). Concordantly, RNAi of *vdac-1*, *F01G4.6*, *spg-7*, *atp-5*, or treatment with the mPTP inhibitor cyclosporine A (CsA) also significantly extends the lifespan of *sgk-1* mutants (Figures 3F–3H, S3D, and S3E). Intestine-specific RNAi of *ant-1.1* in *rict-1* mutants also partially restores normal lifespan, consistent with intestine as a major site of mPTP action in modulation of longevity, as for mTORC2 and autophagy components (Figure S3F).

Given the important role of *sgk-1* in regulation of the mPTP, we examined mitochondrial morphology and function in *sgk-1* mutants. By transmission electron microscopy, mitochondria in *sgk-1* mutants exhibit more fragmentation, more dense morphology, and less apparent cristae (Figure 3I). Oxygen consumption and ATP levels are decreased in mTORC2 mutant worms, as are levels of the electron transport chain protein cytochrome C (Figures 3J, 3K, and S3G). Reduced oxidative capacity of *sgk-1* mutants is not due to decreased mitochondrial mass, as indicated by quantification of mitochondrial DNA and a mitochondrially localized RFP reporter (Figures S3H and S3I), but rather may be due to greater mPTP opening and more fragmented mitochondria. RNAi of electron transport chain genes *nuo-1*, *nuo-6*, and *cco-4* all significantly increase SGK-1 protein levels (Figures S3J and S3K). Together with the observation that mTORC2-SGK1 regulate mitochondrial function, this suggests that SGK-1 is activated by mitochondrial stress as part of a negative feedback loop maintaining mitochondrial homeostasis.

SGK-1 Negatively Regulates mPTP Function by Decreasing VDAC-1 Protein Levels

Levels of VDAC1 are associated with mPTP opening frequency (Tomasello et al., 2009). As SGK1 and VDAC1 physically interact (Figures 3A–3C), it logically follows that SGK-1 may regulate mPTP function by regulating VDAC1 abundance. In order to measure VDAC-1 protein in *sgk-1* mutant worms, we created one integrated and two extrachromosomal transgenic *C. elegans* lines expressing Flag-tagged VDAC-1 (Figures S4A and S4B). VDAC-1-Flag levels are elevated ~2.5-fold in *sgk-1* mutants (Figure 4A), in spite of equivalent mRNA levels (not shown). Native VDAC1 protein level is also higher in *Sgk1* knockout primary hepatocytes (Figure 4B). In contrast, ANT protein in *sgk-1* mutant *C. elegans* and ANT1 and VDAC2 protein levels in *Sgk1* knockout hepatocytes remain unchanged (Figures S4C and S4D). The proteasome inhibitor MG132 eliminated the difference between WT and *Sgk1* knockout hepatocytes (Figure 4C), indicating that SGK1 regulates proteasomal VDAC1 degradation. Expression of constitutively active SGK1(S422D) increases the level of ubiquitinated VDAC1 to a greater extent than WT SGK1, an effect eliminated by the PI3K inhibitor Wortmannin (Figures 4D and 4E). Finally, overexpression of SGK1(S422D) decreases co-expressed VDAC1 protein levels when compared with SGK1(S422A) (Figure 4F), as the latter lacks the capacity to be activated by mTORC2 (Park et al., 1999).




Figure 3. SGK-1 Regulates the mPTP through Physical Interaction with VDAC-1

(A) Purified *C. elegans* Myc-tagged VDAC-1 protein is pulled down by GST-SGK-1 and not GST alone *in vitro*.
 (B) Endogenous VDAC1 is pulled down by native anti-SGK1 coIP.
 (C) Endogenous SGK1 is pulled down by native anti-VDAC1 coIP.
 (D) Calcium retention capacity assay in mouse *Sgk1(-/-)* and WT AML12 cells (arrows indicate calcium pulses). Equal numbers (1×10^7) of cells were used and equivalent biomass was confirmed by western blotting for actin (right).
 (E) RNAi of *ant-1.1* restores normal lifespan in *rict-1* and *sgk-1* mutants (log rank test).
 (F) RNAi of *vdac-1* partially restores normal lifespan in *sgk-1* mutants (log rank test).
 (G) Lifespan extension by RNAi of *F01G4.6* in *sgk-1* mutants and WT *C. elegans* (log rank test).
 (H) RNAi of *spg-7* extends lifespan in *sgk-1* mutants (log rank test).
 (I) Fragmentation and abnormal cristae in *sgk-1* mutants versus WT by electron microscopy (arrows indicate mitochondria). Scale bar, 2 μ m.
 (J) Oxygen consumption is decreased in both *rict-1* and *sgk-1* mutants versus WT (n = 6, one-way ANOVA).
 (K) Western blot for cytochrome C in WT animals and *rict-1* and *sgk-1* mutants (n = 4, one-way ANOVA).
 For tabular survival data and biological replicates see also Table S1. *p < 0.05, **p < 0.01, ****p < 0.0001. All bars indicate means and SEM. See also Figure S3.

We reasoned that direct VDAC1 phosphorylation by SGK1 may prompt VDAC1 degradation. Constitutively active SGK1(S422D) increases serine phosphorylation of mammalian VDAC1 protein *in vitro* (Figure 4G). Mass spectrometric analysis indicates that Ser104 of VDAC1 is a bona fide, direct, but non-canonical SGK1 target (Figure 4H). In confirmation, VDAC1(S104A), which cannot be phosphorylated, reduces baseline serine phosphorylation and eliminates SGK1-mediated VDAC1 serine phosphorylation *in vitro* (Figure 4I). Ubiquitination level of VDAC1(S104A) is decreased versus WT VDAC1 when co-expressed with activated SGK1 (Figure 4J). Further,

VDAC1(S104A) is resistant to SGK1-mediated protein degradation seen with WT VDAC1 protein (Figure 4K). In contrast, mutation of other serine residues on VDAC1 did not prevent SGK1-mediated reductions in protein level (Figures 4K and S4E). Taken together, these data indicate that Ser104 phosphorylation of VDAC1 by SGK1 increases its ubiquitination and subsequent cellular clearance.

Accumulation of VDAC-1 Protein Induces Autophagy

Opening of the mPTP is associated with activation of autophagy (Elmore et al., 2001), and VDAC1 regulates mitochondrial




Figure 4. SGK1 Modulates VDAC1 Protein Ubiquitination and Degradation by Regulating Its Phosphorylation

(A) Western blot for VDAC1::Flag protein levels in *sgk-1* mutant worms versus WT (n = 3, Student's t test).

(B) Western blotting for native VDAC1 protein in primary hepatocytes from *Sgk1^{Lko}* versus *Sgk1^{flox/flox}* control mice (control) (n = 3 cell samples per group, each sample from 1 mouse, Student's t test).

(legend continued on next page)

autophagy (Geisler et al., 2010). Because VDAC-1 is increased in *sgk-1* mutants, we analyzed autophagy with RNAi of *vdac-1*, which inhibits mPTP opening (Tomasello et al., 2009). Knockdown of *vdac-1* in *sgk-1* mutants decreases numbers of GFP::LGG-1 puncta and cleavage of GFP::LGG-1 to free GFP (Figures 5A and 5B). RNAi of *ant-1.1* and treatment with CsA also reduces cleavage of GFP::LGG-1 to free GFP in *sgk-1* mutants (Figures 5C and 5D). *Vdac1* overexpression increases LC3A/B I/II conversion in mammalian cells, indicating that the association of mPTP opening with autophagy is conserved (Figure 5E). Taken together, these data demonstrate that aberrant mPTP opening attributable to excess VDAC-1 increases autophagy levels. Finally, double RNAi of *bec-1* and *ant-1.1* does not additively extend lifespan of *sgk-1* mutants, suggesting action in the same genetic pathway (Figure S5A).

Our earlier result that HLH-30 levels increase in mTORC2 mutants suggests that mTORC2 and VDAC-1 may regulate autophagy at the transcriptional level. mRNA levels of *lgg-1*, *unc-51*, and *hlh-30*, but not *bec-1*, are significantly increased in *rict-1* and *sgk-1* mutants as well as in transgenic *vdac-1* overexpressing animals (Figures 5F and 5G). These data suggest that elevated VDAC-1 levels stimulate autophagy at least partially transcriptionally.

Increased Mitochondrial Permeability Is Detrimental to Longevity and Organismal Health

To explore the impact of increased mitochondrial permeability on aging, we first confirmed that overexpression of both mouse *Vdac1* and *C. elegans vdac-1* in HEK293 cells facilitates mPTP opening (Figure 5H). Second, remarkably, *vdac-1* overexpression alone in *C. elegans* is sufficient to shorten lifespan and decrease cytochrome C protein versus WT animals (Figures 5I, 5J, S5B, and S5C). As hypothesized based upon our observation that activated SGK-1 reduces VDAC-1 protein levels, *sgk-1* overexpression normalizes the shortened lifespan of *vdac-1* transgenics (Figure S5D).

Opening of the mPTP is also triggered by paraquat treatment in mammalian cells (Costantini et al., 1995). As expected, *rict-1* mutants, *sgk-1* mutants, and *vdac-1* overexpression worms, which demonstrate facilitated mPTP opening, exhibit decreased survival with paraquat (Figures S5E and S5F).

Our results suggest that autophagy exerts a progeric, rather than a beneficial effect, in the setting of elevated mitochondrial

permeability. To formally test this, we inhibited autophagy in our *vdac-1* overexpression transgenic worms. Indeed, in *vdac-1* overexpressers, RNAi of *bec-1* restores normal lifespan, as does RNAi of *ant-1.1* (Figures 5K–5L). RNAi of *bec-1* or *ant-1.1* also attenuates paraquat toxicity in *rict-1* and *sgk-1* mutants and *vdac-1* overexpressers, and even moderately in WT animals (Figures S5G–S5K). In aggregate, these results demonstrate that the toxic accumulation of VDAC-1 protein in *sgk-1* mutants is responsible for three major defects: increased mitochondrial permeability, increased autophagy, and shortened lifespan.

One possibility is that blocking autophagy extends lifespan in *sgk-1* mutants by activating the mitochondrial unfolded protein response (via *atfs-1*) (Lamech and Haynes, 2015) or oxidative stress responses (via *FoxO/daf-16* and *Nrf2/skn-1*) (Blackwell et al., 2015). RNAi of *bec-1* extends lifespan in both *atfs-1;sgk-1* and *daf-16;sgk-1* double mutant worms (Figures S6A and S6B), as does double RNAi of *bec-1/skn-1* (Figure S6C). Therefore, the lifespan extending properties of autophagy RNAi in *sgk-1* mutants does not require canonical mitochondrial defenses.

Inhibition of Autophagy Restores Lifespan by Enhancing Mitochondrial Function in *sgk-1* Mutant Worms

We reasoned that excessive clearance of dysfunctional mitochondria could explain the detrimental effects of autophagy on lifespan in the setting of increased mitochondrial permeability. In line with this possibility, RNAi of *bec-1* or *vdac-1* improves oxygen consumption in *sgk-1* mutants (Figures 6A and S6D–S6F). RNAi of *bec-1* also improves ATP levels in *sgk-1* mutants (Figure 6B). In addition, RNAi of *bec-1*, *lgg-1*, and *unc-51* all increase cytochrome C protein levels in *sgk-1* mutants (Figure 6C). Taken together, these data confirm that autophagy has a negative impact on mitochondrial functional capacity when mitochondrial permeability is increased.

We next tested whether the reduction in mitochondrial capacity is causally linked to shortened lifespan in *sgk-1* mutants. To this end, we examined the impact of superimposition of severe mitochondrial electron transport dysfunction in *sgk-1* mutants. In a short-lived mitochondrial complex II mutant *mev-1(kn1)* background (Ishii et al., 1998), RNAi of *bec-1* does not extend the lifespan of *sgk-1* mutants (Figure 6D). Thus, when autophagy inhibition can no longer increase overall mitochondrial oxidative capacity, as in the case of the *mev-1;sgk-1* double mutant, it can no longer restore normal lifespan.

(C) Native VDAC1 level by western blot in control *Sgk1^{flox/flox}* and *Sgk1^{Lko}* primary hepatocytes with or without proteasome inhibitor MG132 (5 μ M for 16 h) (n = 6 cell samples per group, each sample from 1 mouse, Student's t test).

(D and E) Overexpression of active SGK1-S422D (D) increases VDAC1 ubiquitination by western blotting for HA-Ub and (E) is unaffected by the PI3K inhibitor Wortmannin. 293T cells were co-transfected with WT SGK1 (SGK1-WT) or active SGK1 (SGK1-S422D) with VDAC1-Myc and HA-ubiquitin for 48 h. Results shown are representative of 3 biological replicates.

(F) Overexpression of active SGK1-S422D decreases co-transfected VDAC1 protein level versus overexpression of inactive SGK1-S422A (n = 4, Student's t test).

(G) *In vitro* kinase assay shows increased serine phosphorylation of VDAC1 after incubation with SGK1 by western blot (n = 3, Student's t test).

(H) MS2 spectra identifies VDAC1 Ser104 (S¹⁰⁴) as the SGK1 phosphorylation site (n = 2, Student's t test).

(I) SGK1 phosphorylates serine in WT VDAC1 but not VDAC1-S104A *in vitro* (n = 3, two-way ANOVA).

(J) Ubiquitination of VDAC1 and VDAC1-S104A with overexpression of active SGK1-S422D by western blot. Results shown are representative of 3 biological replicates.

(K) Overexpression of active SGK1-S422D decreases WT VDAC1, VDAC1-S101/102A but not VDAC1-S104A protein levels compared with inactive SGK1-S422A overexpression (n = 3, two-way ANOVA).

*p < 0.05, **p < 0.01. All bars indicate means and SEM.

See also Figure S4.




Figure 5. mPTP Opening Regulates Autophagy and Accelerates Aging

(A and B) RNAi of *vdac-1* decreases GFP::LGG-1 and free GFP protein levels (A) and decreases GFP::LGG-1 foci (B) in intestine of *sgk-1* mutants. Scale bar, 20 μm (n = 3, two-way ANOVA for A; n = 60 for ev and n = 51 for *vdac-1*, Student's t test for B). (C and D) In *sgk-1* mutants, RNAi of *ant-1.1* (C) or CsA treatment (D) decreases GFP::LGG-1 or free GFP protein levels (n = 3, two-way ANOVA). (E) Autophagic flux determined by LC3A/B I/II levels detected in 293T cells following overexpression of *Vdac1* for 48 h with and without 50 μM chloroquine (CQ) for 4 h (n = 3, two-way ANOVA). (F and G) *bec-1*, *lgg-1*, *unc-51* (F), and *hlh-30* (G) mRNA levels in L4 stage WT, *ric1-1*, *sgk-1* mutants, and *vdac-1* overexpression transgenics (n = 6, two-way ANOVA [F]; n = 6, one-way ANOVA [G]). (H) mPTP opening by calcium retention assay (arrows indicate calcium pluses) in 293T cells expressing Myc-tagged mouse VDAC1 and *C. elegans* VDAC-1 (A). Equal numbers (2 × 10⁶) of cells were used and confirmed by blotting for tubulin (right). (I) Decreased cytochrome C protein in *vdac-1* transgenic *C. elegans* (n = 3, Student's t test).

(legend continued on next page)

Our results suggest that enhanced mitochondrial clearance may be detrimental to *sgk-1* mutants. Mitochondrial fragmentation has been shown to facilitate mitophagy (Burman et al., 2017). To test whether fragmentation in *sgk-1* mutants (Figure 3I) is responsible for increased mitochondrial clearance and decreased lifespan, we performed RNAi of the mitochondrial fission gene *drp-1*. Knockdown of *drp-1* during larval development significantly extends lifespan in *sgk-1* mutants (Figure 6E). However, knockdown of mitophagy components *pink-1* and *pdr-1/PAKN*, only partially restore normal lifespan (Figures S6G and S6H). Thus, a mitochondrial clearance pathway dependent upon *drp-1* and *bec-1* is principally responsible for decreased *sgk-1* mutant lifespan.

Enhanced Autophagy Is Not Beneficial for Lifespan Extension with Increased Mitochondrial Permeability

Multiple longevity paradigms mechanistically depend upon autophagy, including long-lived *C. elegans* mutants lacking germline stem cells (*glp-1*), calorically restricted *eat-2* mutants, and deficiency of *daf-2/InsulinR* (Hansen et al., 2018). We hypothesize that these paradigms are based upon a foundation of low mitochondrial permeability, as our results suggest that autophagy is transformed into a destructive force whenever mitochondrial permeability is increased. To determine if facilitated mPTP opening is sufficient to mitigate the beneficial effects of increased autophagy, we knocked down a member of mitochondrial complex II, *sdhb-1*, which induces mitochondrial ROS generation and facilitates mPTP opening (Chouchani et al., 2014; Pujol et al., 2013). Consistent with our hypothesis, RNAi of *sdhb-1* eliminated lifespan extension in both *eat-2* and *glp-1* mutants (Figures S6I and S6J). Second, *sgk-1;glp-1* and *sgk-1;eat-2* double mutant worms do not have a long-lived phenotype versus WT, as loss of *sgk-1* similarly facilitates mPTP opening (Figures 6F and 6G). Finally, direct stimulation of mPTP opening by *vdac-1* overexpression in both *eat-2* and *glp-1* mutants completely eliminates lifespan extension seen in these animals (Figures 6H and 6I). Remarkably, *daf-2* knockdown in *vdac-1* overexpression transgenics still extends lifespan (Figure S6K), because *daf-2* deficiency can independently reduce VDAC-1 levels (Figure S6L).

To further extend the generality of these findings, we next examined RNAi of two mitochondrial genes *frh-1* or *nuo-6*, which is known to upregulate autophagy levels, and extend lifespan in an autophagy-dependent manner (Schiavi et al., 2013; Yang and Hekimi, 2010). *vdac-1* overexpression completely suppressed the autophagy-dependent lifespan extension by *frh-1* or *nuo-6* knockdown (Figure 6J). Thus, autophagy only benefits longevity under conditions of low mitochondrial permeability.

Liver-Specific Sgk Knockout Mice Are More Sensitive to Hepatic I/R-Induced Injury

Hepatic ischemia/reperfusion (I/R) injury during liver surgery or liver transplantation negatively influences prognosis. The open-

ing of mPTP in the first few minutes of reperfusion is known to be a critical determinant of I/R injury (Ong et al., 2015). Preventing its opening at this time using mPTP inhibitors such as CsA has been reported to reduce I/R injury in preclinical and clinical studies (Ong et al., 2015).

Given the importance of mPTP opening in I/R injury, we hypothesized that *Sgk1*^{Lko} mice may have increased sensitivity to hepatic I/R injury due to increased mitochondrial permeability and autophagy (Figures 3D, S1E, and S1F). Indeed, following hepatic I/R (Figure 7A), serum alanine transaminase (ALT) and serum aspartate transaminase (AST) levels are more than doubled in *Sgk1*^{Lko} versus control mice (Figures 7B and 7C). Apoptosis (indicated by cleaved Caspase 3) and autophagy (indicated by LC3A/B II) are also higher in *Sgk1*^{Lko} mice after I/R injury (Figure 7D). Concordantly, *Sgk1*^{Lko} mice also have elevated hepatic mRNA levels of pro-inflammatory cytokines interleukin (IL)-1 β and IL-6 (Figure 7E).

In order to address potential functional redundancy in the *Sgk* kinase family, we generated mice lacking *Sgk1*, *Sgk2*, and *Sgk3* in liver (*SgkTKO*). Similar to *Sgk1* knockout hepatocytes, *SgkTKO* hepatocytes demonstrate facilitated mPTP opening in the calcium retention assay (Figure S7A). VDAC1 protein level and the LC3 II/LC3 I ratio are also higher in *SgkTKO* hepatocytes (Figure S7B). To determine whether mPTP opening under I/R is causal in increased hepatic injury in *Sgk* deficiency, we pharmacologically inhibited mPTP opening by pretreatment of *SgkTKO* mice with CsA. Remarkably, mPTP inhibition with CsA reduced post-injury ALT and AST levels in *SgkTKO* mice to a level comparable to control mice (Figures 7F and 7G). Concordantly, levels of cleaved caspase 3 and LC3A/B II protein were restored to control levels in the CsA-treated *SgkTKO* group (Figure 7H).

DISCUSSION

Our results demonstrate a mechanism connecting the nutrient and growth factor responsive kinase SGK1 to health and longevity through its regulation of mitochondrial permeability and autophagy. Work presented here suggests a model whereby the status of the mPTP is a major determinant of the health promoting effects of autophagy (Figure 7I). This represents a significant conceptual advance in our understanding of the way that autophagy modulates health and longevity.

mTORC2 and SGK1 Maintain Mitochondrial Homeostasis

We conclude that mitochondrial dysfunction in the setting of mTORC2 deficiency is related to increased propensity to open the mPTP due to toxic accumulation of VDAC-1 protein. Facilitated mPTP opening in *rict-1* and *sgk-1* mutant worms, in turn, directly compromises oxidative capacity and ATP and

(J) *vdac-1* overexpression shortens lifespan versus WT *C. elegans* (log rank test).

(K and L) RNAi of *bec-1* (K) or *ant-1.1* (L) extends lifespan of the *vdac-1* overexpression strain (log rank test).

For tabular survival data and biological replicates see also Table S1. *p < 0.05, **p < 0.01, ****p < 0.0001. All bars indicate means and SEM. See also Figure S5.




Figure 6. Low Mitochondrial Permeability Is Required for Autophagy-Dependent Lifespan Extension

(A and B) Basal and maximal (FCCP) oxygen consumption (A) and ATP level (B) in *sgk-1* mutants with empty vector (ev) or *bec-1* RNAi (n = 12, two-way ANOVA; n = 7, Student's t test).

(C) Cytochrome C western blot in *sgk-1* mutants following RNAi of *bec-1*, *lgg-1*, or *unc-51* (n = 4, one-way ANOVA).

(D) *bec-1* RNAi does not extend lifespan in *mev-1(kn1);sgk-1* double mutants.

(E) Larval knockdown of *drp-1* extends lifespan in *sgk-1* mutant worms (log rank test).

(F–I) Elimination of autophagy-dependent lifespan extension in both *eat-2(ad465)* mutants with (F) *sgk-1* loss of function and (H) *vdac-1* overexpression and in *glp-1(e2141)* mutants with (G) *sgk-1* loss of function and (I) *vdac-1* overexpression (log rank test).

(J) RNAi of *frh-1* and *nuo-6* extends lifespan in WT worms but has no effect on *vdac-1* transgenic worms (log rank test).

For tabular survival data and biological replicates see also Table S1. *p < 0.05, **p < 0.01, ****p < 0.0001. All bars indicate means and SEM.

See also Figure S6.

cytochrome C protein levels. Thus, progeria in *C. elegans* mTORC2 pathway mutants is a direct consequence of increased mPTP opening and ensuing mitochondrial dysfunction.

mTORC2 is known to regulate mitochondrial function by maintaining mitochondria-associated endoplasmic reticulum (ER) membrane integrity (Betz et al., 2013). Our data demonstrate a previously unappreciated dimension of mitochondrial regulation by mTORC2. SGK1 has been reported to localize to mitochon-

dria under stress, which permits access to physiologically appropriate mitochondrial interacting proteins and substrates, as part of cellular stress responses (O'Keefe et al., 2013). We conclude that a major way that mTORC2 and SGK1 regulate mitochondrial permeability is through physical association of mTORC2 with mPTP regulatory molecules, modulating the abundance of the outer mitochondrial membrane protein VDAC1 through SGK1.




Figure 7. Mice Lacking *Sgk* Are More Sensitive to Liver I/R Injury

(A) Hepatic warm I/R injury model.

(B and C) Serum ALT (B) and AST (C) levels in 12-week-old *Sgk1^{Lko}* versus control *Sgk1^{fllox/fllox}* mice after hepatic I/R injury (n = 10 per group, Student's t test).

(D) Active, cleaved caspase 3 and LC3A/B II protein levels mice after hepatic I/R injury (n = 5 for control and n = 6 for *Sgk1^{Lko}*, Student's t test).

(E) mRNA levels of inflammatory genes IL-1 β and IL-6 by qPCR (n = 11 per group, Student's t test).

(F and G) Serum ALT (F) and AST (G) levels of 24-week-old *SgkTKO* mice versus *Sgk1^{fllox/fllox}* control after hepatic I/R injury with or without CsA pretreatment (n = 8 for control and *SgkTKO*, and n = 7 for *SgkTKO*+CsA, one-way ANOVA).

(H) Active, cleaved caspase 3 and LC3A/B protein levels in liver after hepatic I/R injury with or without CsA pretreatment (n = 7 for control and *SgkTKO*+CsA, and n = 8 for *SgkTKO*, two-way ANOVA).

(I) Model of the effect of autophagy on lifespan with facilitated opening of the mPTP (mPTP open) versus normal mitochondrial permeability (mPTP closed) conditions.

*p < 0.05, **p < 0.01. All bars indicate means and SEM.

mTORC2 and SGK-1 Suppress Maladaptive Autophagy

We suggest that increased intestinal autophagic flux in *sgk-1* mutant worms has negative consequences for longevity. However, our data leave open the possibility that autophagy is important in additional tissues in modulation of longevity downstream of mTORC2. Autophagic activity is regulated spatiotemporally in multiple tissues during aging (Chang et al., 2017). Spatiotemporal examination of autophagy in

rict-1 and *sgk-1* mutants may provide greater mechanistic understanding of how autophagy contributes to progeria. Finally, while our data suggest cell autonomous modulation of the mPTP and autophagy by mTORC2, they do not exclude the possibility of non-cell autonomous regulation of autophagy.

Data presented here suggest that a mechanism by which mPTP opening triggers autophagy is through transcriptional

induction of the key autophagy transcription factor TFEB/HLH-30. Opening of the mPTP also induces ROS production (Batandier et al., 2004), and two major ROS H_2O_2 and O_2^- have been suggested to regulate autophagy (Scherz-Shouval and Elazar, 2011). Further, VDACs specifically interact with Parkin on defective mitochondria, targeting mitochondria for subsequent mitophagy (Geisler et al., 2010). Future work is needed to determine precisely how mPTP opening leads to induction of autophagy.

While mTORC2 regulates autophagy through modulation of the mPTP, our work leaves open the possibility of additional mechanisms. mTORC2-SGK signaling has been shown to activate mTORC1 (Castel et al., 2016), and mTORC1 plays a central role in suppression of autophagy (Saxton and Sabatini, 2017). Therefore, mTORC2 may indirectly reduce autophagy through SGK by activating mTORC1. Consistent with this possibility, our results show decreased mTORC1 activity in *SgkTKO* hepatocytes (Figure S7C). Irrespective of the mechanism, we conclude that a major role of mTORC2 and SGK1 is to act as autophagy gatekeepers.

mPTP Opening Negatively Modulates Longevity and Health

The work presented here indicates that in the absence of functional mTORC2 or SGK-1, increased opening of the mPTP drives both mitochondrial dysfunction and increased autophagy. We put forward that this combination is particularly devastating, shortening lifespan and increasing susceptibility to I/R injury.

Our results mechanistically connect increased mPTP opening to shortened lifespan and hypersensitivity to oxidative stress. Emerging data indicate that mitochondrial permeability increases in aged animals and humans and in aging-related degenerative diseases, and diverse manipulations that promote health and extend lifespan decrease mPTP opening (Rottenberg and Hoek, 2017). Caloric restriction protects mitochondria from mPTP opening in rat heart (Hofer et al., 2009). Metformin, an antidiabetic drug with anti-aging properties, inhibits mPTP opening (Bhamra et al., 2008). Thus, maintenance of low mitochondrial permeability may be an obligate step in order to extend lifespan.

Autophagy as a Double-Edged Sword in I/R Injury

The role of autophagy in I/R injury is complex. Beneficial effects of autophagy during I/R have been attributed to ATP generation, improved cellular homeostasis and decreased expression of inflammatory genes (Ma et al., 2015). On the other hand, excessive autophagy contributes to pressure overload-induced cardiac remodeling and heart failure in cardiac I/R injury (Gottlieb and Mentzer, 2010). In a rat hepatic I/R injury model, chloroquine, which inhibits autophagic flux, prevents ischemic liver damage at the early phase, but aggravates liver damage at the late phase in liver I/R injury (Fang et al., 2013).

Based upon our results, we hypothesize that the status of the mPTP is a major governing influence on the effects of autophagy on ischemia-reperfusion injury. Because widespread mPTP opening occurs shortly after reperfusion (Ong et al., 2015), and the ensuing mitochondrial permeability drives autophagy to be detrimental, inhibition of autophagy during early re-

perfusion reduces injury. At the late phase of reperfusion, the mPTP closes in mitochondria that are able to restore functionality, and autophagy plays a beneficial role by selectively degrading dysfunctional mitochondria and improving cellular homeostasis (Ma et al., 2015). Recent work suggests that accumulation of succinate is responsible for mitochondrial ROS production during I/R injury (Chouchani et al., 2014), sensitizing mitochondria to mPTP opening (Briston et al., 2017). It is tantalizing to speculate that mTORC2 mutants are more sensitive to succinate-induced mPTP opening because of accumulation of VDAC1.

Resolving the Paradox: How Inhibition of Autophagy Restores Lifespan in mTORC2 Mutants

Increased autophagy is generally thought to be beneficial for longevity, as almost every manipulation that extends lifespan is dependent on autophagy (Madeo et al., 2015). Enhancing autophagy using genetic methods by activating Beclin increases longevity in mice, and similarly, overexpression of *hlf-30* in *C. elegans* also extends lifespan (Fernández et al., 2018; Lapierre et al., 2013). However, we conclusively demonstrate that elevated autophagy is detrimental for organismal health when coupled with increased mitochondrial permeability. We surmise that in mTORC2 mutant worms, increased mitochondrial permeability triggers mitochondrial fragmentation and initiates autophagy, resulting in excessive, non-selective mitochondrial clearance and failure to replace those mitochondria with normal, healthy organelles. Thus, inhibition of autophagy in mTORC2 mutants restores normal lifespan by increasing mitochondrial mass and restoring oxidative capacity. This is further supported by the failure of lifespan extension under *bec-1* RNAi in *sgk-1/mev-1* double mutants, indicating that mitochondria with intact electron transport chain activity are required to restore normal lifespan.

Several lines of evidence support the idea that increased mitochondrial permeability rather than a defect in ATP synthase activity underlies lifespan shortening in mTORC2 mutants. First, RNAi of ATP synthase extends lifespan (Dillin et al., 2002; Lee et al., 2003). Second, knockdown of *ant-1.1* or *atp-5*, which decreases ATP production, further extends lifespan in *sgk-1* mutant worms. Finally, RNAi of *bec-1* or *vdac-1* increases oxygen consumption and ATP level in *sgk-1* mutant worms but not in WT worms, suggesting the energetic defect in these mutants is due to mPTP dysregulation and ensuing autophagy rather than a primary deficit in energy generation.

The work presented here suggests an over-arching idea that cellular context determines whether autophagy has beneficial or detrimental effects on health. An emerging theme in longevity studies is that in order for a manipulation to extend lifespan, it must suppress mitochondrial permeability. Further, increases in mitochondrial permeability accompany aging and many disease states, and our work indicates that these increases in mitochondrial permeability are causal in aging. Our data also intimates that autophagy induced during I/R injury is harmful, exacerbating damage, if accompanied by increases in mPTP opening. Future work will be needed to determine whether mitochondrial permeability can be favorably manipulated in

preclinical and clinical studies in order to promote healthy aging and reduce morbidity associated with diseases such as ischemic stroke, heart attack, ischemic renal and hepatic disease, and with organ transplantation.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **CONTACT FOR REAGENT AND RESOURCE SHARING**
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Nematode Strains
 - *E. coli* Strains
 - Mice
 - Cell Lines, Culture Conditions and Transfection
- **METHOD DETAILS**
 - Generation of Sgk1 Knockout AML12 Cells
 - Transmission Electron Microscopy (TEM)
 - Measuring Autophagy Levels in *C. elegans* and Cells
 - Chloroquine *C. elegans* Treatment
 - Western Blotting
 - Longevity Assay
 - RNA Interference (RNAi)
 - Quantitative RT-PCR
 - Brood Size
 - Developmental Timing
 - Nile-Red Staining
 - SGK-1-GFP Co-IP for Mass Spectrometry
 - Oxygen Consumption
 - ATP Measurement
 - Creation of the VDAC-1::FLAG *C. elegans* Transgenic Lines
 - VDAC1 In Vitro Pull-Down Assay and Ubiquitination Assay
 - mtDNA Copy Number
 - Mitochondrial Ca²⁺ Uptake
 - Paraquat Treatment
 - Identification of the VDAC1 Phosphorylation Site by Mass Spectrometry
 - Hepatic Ischemia/Reperfusion Model
- **QUANTIFICATION AND STATISTICAL ANALYSIS**

SUPPLEMENTAL INFORMATION

Supplemental Information can be found with this article online at <https://doi.org/10.1016/j.cell.2019.02.013>.

ACKNOWLEDGMENTS

We thank Joseph Avruch, Noriko Oshiro-Rapley, Ning Dai, and Laura Regue for discussions and reagents; Johnathan Whetstone for the HA-Ub plasmid; and Diane Capen at the Program in Membrane Biology at MGH for EM. Thanks to the NIH/NIDDK-funded NORC of Harvard (P30DK040561) and the NIH/NIDDK-funded Boston-Area DERC (P30DK057521) for core services. Some strains were provided by the CGC, funded by the NIH Office of Research Infrastructure Programs (P40OD010440), and the *C. elegans* Knockout Consortium.

This work was funded by NIH (K08DK087941 and R01DK101522 to A.A.S., R01AG038664 to M.H., and R21AG058038 to C.K.), the Ellison Medical Foundation New Scholar in Aging Award (to A.A.S.), the Howard M. Goodman Fellowship (to A.A.S.), the Charles H. Hood Foundation Child Health Research Award (to A.A.S.), the Weissman Family MGH Research Scholar Award (to A.A.S.), a Glenn Award for Research in Biological Mechanisms of Aging (to A.A.S.), the Julie Martin Award from The Ellison Medical Foundation/AFAR (to M.H.), and an MGH ECOR Postdoctoral Fellowship (to B.Z.).

AUTHOR CONTRIBUTIONS

Conceptualization, B.Z. and A.A.S.; Methodology, B.Z., L.W., K.J.K., S.D., and A.A.S.; Formal Analysis, B.Z., C.K., J.K., M.H., W.H., S.D., and A.A.S.; Investigation, B.Z., C.K., J.K., L.W., K.J.K., M.C.K., C.M.W., L.C., Y.Z., S.L., and A.A.S.; Resources, C.K., M.H., G.F.-T., and A.N.-F.-T.; Writing – Original Draft, B.Z. and A.A.S.; Writing – Review & Editing, B.Z., M.C.K., M.H., and A.A.S.; Visualization, B.Z., M.C.K., and A.A.S.; Supervision, A.A.S.; Funding Acquisition, A.A.S.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: April 4, 2018

Revised: November 21, 2018

Accepted: February 11, 2019

Published: March 28, 2019

REFERENCES

- Batandier, C., Leverage, X., and Fontaine, E. (2004). Opening of the mitochondrial permeability transition pore induces reactive oxygen species production at the level of the respiratory chain complex I. *J. Biol. Chem.* 279, 17197–17204.
- Betz, C., Stracka, D., Prescianotto-Baschong, C., Frieden, M., Demareux, N., and Hall, M.N. (2013). Feature Article: mTOR complex 2-Akt signaling at mitochondria-associated endoplasmic reticulum membranes (MAM) regulates mitochondrial physiology. *Proc. Natl. Acad. Sci. USA* 110, 12526–12534.
- Bhamra, G.S., Hausenloy, D.J., Davidson, S.M., Carr, R.D., Paiva, M., Wynne, A.M., Mocanu, M.M., and Yellon, D.M. (2008). Metformin protects the ischemic heart by the Akt-mediated inhibition of mitochondrial permeability transition pore opening. *Basic Res. Cardiol.* 103, 274–284.
- Blackwell, T.K., Steinbaugh, M.J., Hourihan, J.M., Ewald, C.Y., and Isik, M. (2015). SKN-1/Nrf, stress responses, and aging in *Caenorhabditis elegans*. *Free Radic. Biol. Med.* 88 (Pt B), 290–301.
- Briston, T., Roberts, M., Lewis, S., Powney, B., Staddon, J.M., Szabadkai, G., and Duchon, M.R. (2017). Mitochondrial permeability transition pore: sensitivity to opening and mechanistic dependence on substrate availability. *Sci. Rep.* 7, 10492.
- Buisson, R., Niraj, J., Rodrigue, A., Ho, C.K., Kreuzer, J., Foo, T.K., Hardy, E.J., Dellaire, G., Haas, W., Xia, B., et al. (2017). Coupling of Homologous Recombination and the Checkpoint by ATR. *Mol. Cell* 65, 336–346.
- Burman, J.L., Pickles, S., Wang, C., Sekine, S., Vargas, J.N.S., Zhang, Z., Youle, A.M., Nezhich, C.L., Wu, X., Hammer, J.A., and Youle, R.J. (2017). Mitochondrial fission facilitates the selective mitophagy of protein aggregates. *J. Cell Biol.* 216, 3231–3247.
- Calvo, S.E., Clauser, K.R., and Mootha, V.K. (2016). MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Res.* 44 (D1), D1251–D1257.
- Castel, P., Ellis, H., Bago, R., Toska, E., Razavi, P., Carmona, F.J., Kannan, S., Verma, C.S., Dickler, M., Chandrapatya, S., et al. (2016). PDK1-SGK1 Signaling Sustains AKT-Independent mTORC1 Activation and Confers Resistance to PI3K α Inhibition. *Cancer Cell* 30, 229–242.

- Chang, J.T., Kumsta, C., Hellman, A.B., Adams, L.M., and Hansen, M. (2017). Spatiotemporal regulation of autophagy during *Caenorhabditis elegans* aging. *eLife* 6, e18459.
- Chouchani, E.T., Pell, V.R., Gaude, E., Aksentijević, D., Sundier, S.Y., Robb, E.L., Logan, A., Nadtochiy, S.M., Ord, E.N.J., Smith, A.C., et al. (2014). Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 515, 431–435.
- Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., and Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819–823.
- Cordas, E., Náráy-Fejes-Tóth, A., and Fejes-Tóth, G. (2007). Subcellular location of serum- and glucocorticoid-induced kinase-1 in renal and mammary epithelial cells. *Am. J. Physiol. Cell Physiol.* 292, C1971–C1981.
- Costantini, P., Petronilli, V., Colonna, R., and Bernardi, P. (1995). On the effects of paraquat on isolated mitochondria. Evidence that paraquat causes opening of the cyclosporin A-sensitive permeability transition pore synergistically with nitric oxide. *Toxicology* 99, 77–88.
- Crawford, D., Libina, N., and Kenyon, C. (2007). *Caenorhabditis elegans* integrates food and reproductive signals in lifespan determination. *Aging Cell* 6, 715–721.
- Di Cristofano, A. (2017). SGK1: The Dark Side of PI3K Signaling. *Curr. Top. Dev. Biol.* 123, 49–71.
- Dillin, A., Hsu, A.L., Arantes-Oliveira, N., Lehrer-Graier, J., Hsin, H., Fraser, A.G., Kamath, R.S., Ahringer, J., and Kenyon, C. (2002). Rates of behavior and aging specified by mitochondrial function during development. *Science* 298, 2398–2401.
- Elmore, S.P., Qian, T., Grissom, S.F., and Lemasters, J.J. (2001). The mitochondrial permeability transition initiates autophagy in rat hepatocytes. *FASEB J.* 15, 2286–2287.
- Fang, H., Liu, A., Dahmen, U., and Dirsch, O. (2013). Dual role of chloroquine in liver ischemia reperfusion injury: reduction of liver damage in early phase, but aggravation in late phase. *Cell Death Dis.* 4, e694.
- Fejes-Tóth, G., Frindt, G., Náráy-Fejes-Tóth, A., and Palmer, L.G. (2008). Epithelial Na⁺ channel activation and processing in mice lacking SGK1. *Am. J. Physiol. Renal Physiol.* 294, F1298–F1305.
- Fernández, A.F., Sebti, S., Wei, Y., Zou, Z., Shi, M., McMillan, K.L., He, C., Ting, T., Liu, Y., Chiang, W.C., et al. (2018). Disruption of the beclin 1-BCL2 autophagy regulatory complex promotes longevity in mice. *Nature* 558, 136–140.
- García-Martínez, J.M., and Alessi, D.R. (2008). mTOR complex 2 (mTORC2) controls hydrophobic motif phosphorylation and activation of serum- and glucocorticoid-induced protein kinase 1 (SGK1). *Biochem. J.* 416, 375–385.
- Geisler, S., Holmström, K.M., Skujat, D., Fiesel, F.C., Rothfuss, O.C., Kahle, P.J., and Springer, W. (2010). PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat. Cell Biol.* 12, 119–131.
- Gottlieb, R.A., and Mentzer, R.M. (2010). Autophagy during cardiac stress: joys and frustrations of autophagy. *Annu. Rev. Physiol.* 72, 45–59.
- Han, S.K., Lee, D., Lee, H., Kim, D., Son, H.G., Yang, J.S., Lee, S.V., and Kim, S. (2016). OASIS 2: online application for survival analysis 2 with features for the analysis of maximal lifespan and healthspan in aging research. *Oncotarget* 7, 56147–56152.
- Hansen, M., Rubinsztein, D.C., and Walker, D.W. (2018). Autophagy as a promoter of longevity: insights from model organisms. *Nat. Rev. Mol. Cell Biol.* 19, 579–593.
- Hofer, T., Servais, S., Seo, A.Y., Marzetti, E., Hiona, A., Upadhyay, S.J., Wohlgemuth, S.E., and Leeuwenburgh, C. (2009). Bioenergetics and permeability transition pore opening in heart subsarcolemmal and intermyofibrillar mitochondria: effects of aging and lifelong calorie restriction. *Mech. Ageing Dev.* 130, 297–307.
- Ishii, N., Fujii, M., Hartman, P.S., Tsuda, M., Yasuda, K., Senoo-Matsuda, N., Yanase, S., Ayusawa, D., and Suzuki, K. (1998). A mutation in succinate dehydrogenase cytochrome b causes oxidative stress and ageing in nematodes. *Nature* 394, 694–697.
- Kang, C., You, Y.J., and Avery, L. (2007). Dual roles of autophagy in the survival of *Caenorhabditis elegans* during starvation. *Genes Dev.* 21, 2161–2171.
- Klionsky, D.J., Abdelmohsen, K., Abe, A., Abedin, M.J., Abeliovich, H., Acevedo Arozena, A., Adachi, H., Adams, C.M., Adams, P.D., Adeli, K., et al. (2016). Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy* 12, 1–222.
- Kwong, J.Q., and Molkentin, J.D. (2015). Physiological and pathological roles of the mitochondrial permeability transition pore in the heart. *Cell Metab.* 21, 206–214.
- Lamech, L.T., and Haynes, C.M. (2015). The unpredictability of prolonged activation of stress response pathways. *J. Cell Biol.* 209, 781–787.
- Lapierre, L.R., De Magalhaes Filho, C.D., McQuary, P.R., Chu, C.C., Visvikis, O., Chang, J.T., Gelino, S., Ong, B., Davis, A.E., Irazoqui, J.E., et al. (2013). The TFEB orthologue HLH-30 regulates autophagy and modulates longevity in *Caenorhabditis elegans*. *Nat. Commun.* 4, 2267.
- Lee, S.S., Lee, R.Y., Fraser, A.G., Kamath, R.S., Ahringer, J., and Ruvkun, G. (2003). A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat. Genet.* 33, 40–48.
- Li, W.C., Ralphs, K.L., and Tosh, D. (2010). Isolation and culture of adult mouse hepatocytes. *Methods Mol. Biol.* 633, 185–196.
- López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The hallmarks of aging. *Cell* 153, 1194–1217.
- Ma, S., Wang, Y., Chen, Y., and Cao, F. (2015). The role of the autophagy in myocardial ischemia/reperfusion injury. *Biochim. Biophys. Acta* 1852, 271–276.
- Madeo, F., Zimmermann, A., Maiuri, M.C., and Kroemer, G. (2015). Essential role for autophagy in life span extension. *J. Clin. Invest.* 125, 85–93.
- Manil-Ségalen, M., Lefebvre, C., Jenzer, C., Trichet, M., Boulogne, C., Satiat-Jeuemaitre, B., and Legouis, R. (2014). The *C. elegans* LC3 acts downstream of GABARAP to degrade autophagosomes by interacting with the HOPS subunit VPS39. *Dev. Cell* 28, 43–55.
- Matecic, M., Smith, D.L., Pan, X., Maqani, N., Bekiranov, S., Boeke, J.D., and Smith, J.S. (2010). A microarray-based genetic screen for yeast chronological aging factors. *PLoS Genet.* 6, e1000921.
- Mizunuma, M., Neumann-Haefelin, E., Moroz, N., Li, Y., and Blackwell, T.K. (2014). mTORC2-SGK-1 acts in two environmentally responsive pathways with opposing effects on longevity. *Aging Cell* 13, 869–878.
- O’Keeffe, B.A., Cilia, S., Maiyar, A.C., Vaysberg, M., and Firestone, G.L. (2013). The serum- and glucocorticoid-induced protein kinase-1 (Sgk-1) mitochondria connection: identification of the IF-1 inhibitor of the F(1)F(0)-ATPase as a mitochondria-specific binding target and the stress-induced mitochondrial localization of endogenous Sgk-1. *Biochimie* 95, 1258–1265.
- Ong, S.B., Samangouei, P., Kalkhoran, S.B., and Hausenloy, D.J. (2015). The mitochondrial permeability transition pore and its role in myocardial ischemia reperfusion injury. *J. Mol. Cell. Cardiol.* 78, 23–34.
- Palikaras, K., and Tavernarakis, N. (2016). Intracellular Assessment of ATP Levels in *Caenorhabditis elegans*. *Biol. Protoc.* 6, e22048.
- Park, J., Leong, M.L., Buse, P., Maiyar, A.C., Firestone, G.L., and Hemmings, B.A. (1999). Serum and glucocorticoid-inducible kinase (SGK) is a target of the PI 3-kinase-stimulated signaling pathway. *EMBO J.* 18, 3024–3033.
- Pujol, C., Bratic-Hench, I., Sumakovic, M., Hench, J., Mourier, A., Baumann, L., Pavlenko, V., and Trifunovic, A. (2013). Succinate dehydrogenase upregulation destabilize complex I and limits the lifespan of gas-1 mutant. *PLoS ONE* 8, e59493.
- Rottenberg, H., and Hoek, J.B. (2017). The path from mitochondrial ROS to aging runs through the mitochondrial permeability transition pore. *Aging Cell* 16, 943–955.
- Saxton, R.A., and Sabatini, D.M. (2017). mTOR Signaling in Growth, Metabolism, and Disease. *Cell* 169, 361–371.
- Scherz-Shouval, R., and Elazar, Z. (2011). Regulation of autophagy by ROS: physiology and pathology. *Trends Biochem. Sci.* 36, 30–38.

- Schiavi, A., Torgovnick, A., Kell, A., Megalou, E., Castelein, N., Guccini, I., Marzocchella, L., Gelino, S., Hansen, M., Malisan, F., et al. (2013). Autophagy induction extends lifespan and reduces lipid content in response to frataxin silencing in *C. elegans*. *Exp. Gerontol.* *48*, 191–201.
- Schwartz, D., and Gygi, S.P. (2005). An iterative statistical approach to the identification of protein phosphorylation motifs from large-scale data sets. *Nat. Biotechnol.* *23*, 1391–1398.
- Shanmughapriya, S., Rajan, S., Hoffman, N.E., Higgins, A.M., Tomar, D., Nemani, N., Hines, K.J., Smith, D.J., Eguchi, A., Vallem, S., et al. (2015). SPG7 Is an Essential and Conserved Component of the Mitochondrial Permeability Transition Pore. *Mol. Cell* *60*, 47–62.
- Soukas, A.A., Kane, E.A., Carr, C.E., Melo, J.A., and Ruvkun, G. (2009). Rictor/TORC2 regulates fat metabolism, feeding, growth, and life span in *Caenorhabditis elegans*. *Genes Dev.* *23*, 496–511.
- Thorburn, A. (2014). Autophagy and its effects: making sense of double-edged swords. *PLoS Biol.* *12*, e1001967.
- Tomasello, F., Messina, A., Lartigue, L., Schembri, L., Medina, C., Reina, S., Thoraval, D., Crouzet, M., Ichas, F., De Pinto, V., and De Giorgi, F. (2009). Outer membrane VDAC1 controls permeability transition of the inner mitochondrial membrane in cellulose during stress-induced apoptosis. *Cell Res.* *19*, 1363–1376.
- Wang, M.C., Oakley, H.D., Carr, C.E., Sowa, J.N., and Ruvkun, G. (2014). Gene pathways that delay *Caenorhabditis elegans* reproductive senescence. *PLoS Genet.* *10*, e1004752.
- Yang, W., and Hekimi, S. (2010). Two modes of mitochondrial dysfunction lead independently to lifespan extension in *Caenorhabditis elegans*. *Aging Cell* *9*, 433–447.
- Zhang, H., Chang, J.T., Guo, B., Hansen, M., Jia, K., Kovács, A.L., Kumsta, C., Lapiere, L.R., Legouis, R., Lin, L., et al. (2015). Guidelines for monitoring autophagy in *Caenorhabditis elegans*. *Autophagy* *11*, 9–27.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-GFP	Thermo Fisher Scientific	Cat# MA5-15256; RRID:AB_10979281
Mouse monoclonal anti-HA (clone GT4810)	Sigma-Aldrich	Cat# SAB2702217; RRID:AB_2750919
Mouse monoclonal anti-Flag (M2)	Sigma-Aldrich	Cat# F3165; RRID:AB_259529
Mouse monoclonal anti-Actin (C4)	Abcam	Cat# ab14128; RRID:AB_300931
Mouse monoclonal anti-Cytochrome C (37BA11)	Abcam	Cat# ab110325; RRID:AB_10864775
Mouse monoclonal anti-Myc (9E10)	Abcam	Cat# ab32; RRID:AB_303599
Mouse monoclonal anti-ANT 1 [5F51BB5AG7]	Abcam	Cat# ab110322; RRID:AB_10862212
Rabbit monoclonal anti-Myc (71D10)	Cell Signaling Technology	Cat# 2276; RRID:AB_331783
Rabbit monoclonal anti-LC3A/B (D3U4C)	Cell Signaling Technology	Cat# 12741; RRID:AB_2617131
Rabbit monoclonal anti-SGK1 (D27C11)	Cell Signaling Technology	Cat# 12103; RRID:AB_2687476
Rabbit monoclonal anti-SGK2 (D7G1)	Cell Signaling Technology	Cat# 7499; RRID:AB_10828732
Rabbit monoclonal anti-SGK3 (D18D1)	Cell Signaling Technology	Cat# 8156; RRID:AB_10949507
Rabbit polyclonal anti-VDAC2	Cell Signaling Technology	Cat# 9412; RRID:AB_10828727
Rabbit monoclonal anti-HSP90 (C45G5)	Cell Signaling Technology	Cat# 4877; RRID:AB_2233307
Rabbit monoclonal anti-p70 S6 Kinase (49D7)	Cell Signaling Technology	Cat# 2708; RRID:AB_390722
Rabbit monoclonal anti-Phospho-p70 S6 Kinase (Thr389) (108D2)	Cell Signaling Technology	Cat# 9234; RRID:AB_2269803
Rabbit monoclonal anti-S6 Ribosomal Protein (5G10)	Cell Signaling Technology	Cat# 2217; RRID:AB_331355
Rabbit monoclonal anti-Phospho-S6 Ribosomal Protein (Ser240/244) (D68F8)	Cell Signaling Technology	Cat# 5364; RRID:AB_10694233
Rabbit monoclonal anti-Phospho-AMPK α (Thr172) (40H9)	Cell Signaling Technology	Cat# 2535; RRID:AB_331250
Rabbit monoclonal anti-VDAC (D73D12)	Cell Signaling Technology	Cat# 4661; RRID:AB_10557420
Rabbit monoclonal anti-Caspase-3 (8G10)	Cell Signaling Technology	Cat# 9665; RRID:AB_2069872
Mouse monoclonal anti-Tubulin	Cell Signaling Technology	Cat# 3873; RRID:AB_1904178
Mouse monoclonal anti-GST	Cell Signaling Technology	Cat# 2624; RRID:AB_2189875
Mouse monoclonal anti-VDAC1 (B-6)	Santa Cruz Biotechnology	Cat# sc-390996; RRID:AB_2750920
Mouse monoclonal anti-phospho-Serine (16B4)	Santa Cruz Biotechnology	Cat# sc-81514; RRID:AB_1128624
Bacterial and Virus Strains		
<i>Escherichia coli</i> OP50	Caenorhabditis Genetics Center (CGC)	RRID:WB-STRAIN:OP50
RNAi feeding bacterial strain HT115(DE3)	Source BioScience (Ahringer library)	N/A
RNAi feeding bacterial strain HT115(DE3)	Source BioScience (Vidal library)	N/A
Biological Samples		
Mouse liver tissue from control, <i>Sgk1</i> ^{L^{ko}} , <i>SgkTKO</i> mice	This paper	N/A
Chemicals, Peptides, and Recombinant Proteins		
Cyclosporine A	Enzo Life Sciences	BML-A195
MG132	Enzo Life Sciences	BML-PI102
Paraquat	Sigma-Aldrich	36541
FCCP	Sigma-Aldrich	C2920
Chloroquine	Sigma-Aldrich	C6628
Wortmannin	Sigma-Aldrich	W1628
Floxuridine (FUDR)	Sigma-Aldrich	1271008
Bafilomycin A	BioVotica, Germany	88899-55-2

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Calcium Green-5N	Thermo Fisher Scientific	C3737
RNAzol RT	Molecular Research Center	RN 190
Critical Commercial Assays		
Pierce GST Spin Purification Kit	Thermo Fisher Scientific	16106
Pierce HisPur Ni-NTA Purification Kit	Thermo Fisher Scientific	88227
ATP Determination Kit	Thermo Fisher Scientific	A22066
CYTO-ID Autophagy detection kit 2.0	Enzo Life Sciences	ENZ-KIT175
Experimental Models: Cell Lines		
Human: HEK293T	Gift from Joseph Avruch's lab	Not authenticated
Mouse: AML12	ATCC	CRL-2254
Mouse: Primary hepatocytes	This paper	N/A
Experimental Models: Organisms/Strains		
<i>C. elegans</i> : N2 Bristol	Caenorhabditis Genetics Center	RRID:WB-STRAIN:N2_(ancestral)
<i>C. elegans</i> : TK22 <i>mev-1(kn1)</i> 5x	Caenorhabditis Genetics Center	RRID:WB-STRAIN:TK22
<i>C. elegans</i> : DA465 <i>eat-2(ad465)</i>	Caenorhabditis Genetics Center	RRID:WB-STRAIN:DA465
<i>C. elegans</i> : CB4037 <i>glp-1(e2141)</i>	Caenorhabditis Genetics Center	RRID:WB-STRAIN:CB4037
<i>C. elegans</i> : DA2123 <i>adls2122 [lgg-1p::GFP::lgg-1 + rol-6(su1006)]</i>	Caenorhabditis Genetics Center	RRID:WB-STRAIN:DA2123
<i>C. elegans</i> : RD202 <i>unc-119(ed3)</i> III; <i>ppls??[unc-119(+); lgg-1p::gfp::lgg-1(G116A)]</i>	Legouis lab	N/A
<i>C. elegans</i> : JIN1821 <i>JinIs[h1h-30p::HLH-30::GFP]</i>	This study	N/A
<i>C. elegans</i> : MGH12 <i>mglS60[sgk-1::GFP pRF4]</i>	This study	N/A
<i>C. elegans</i> : MGH48 <i>mglS43[ges-1p::GFP_PTS1]</i>	This study	N/A
<i>C. elegans</i> : MGH55 <i>daf-16(mgDf47)</i> 3x	This study	N/A
<i>C. elegans</i> : MGH361 <i>atfs-1(tm4525)</i> 3x	This study	N/A
<i>C. elegans</i> : MGH266 <i>riect-1(mg451)</i> 8x	This study	N/A
<i>C. elegans</i> : MGH300 <i>sgk-1(mg455)</i> 8x	This study	N/A
<i>C. elegans</i> : MGH358 <i>mev-1(kn1);sgk-1(mg455)</i>	This study	N/A
<i>C. elegans</i> : MGH359 <i>atfs-1(tm4525);sgk-1(mg455)</i>	This study	N/A
<i>C. elegans</i> : MGH360 <i>daf-16(mgDf47);sgk-1(mg455)</i>	This study	N/A
<i>C. elegans</i> : MGH84 <i>mglS54[myo-3p::mito::mRFP3]</i>	This study	N/A
<i>C. elegans</i> : MGH353 <i>sgk-1(mg455);mglS54[myo-3p::mito::mRFP3]</i>	This study	N/A
<i>C. elegans</i> : MGH354 <i>alxEx91[vdac-1p::VDAC-1::FLAG::SL2::GFP] (vdac-1 oe1)</i>	This study	N/A
<i>C. elegans</i> : MGH355 <i>alxEx92[vdac-1p::VDAC-1::FLAG; myo-2p::GFP] (vdac-1 oe2)</i>	This study	N/A
<i>C. elegans</i> : MGH356 <i>alxS28[vdac-1p::VDAC-1::FLAG; myo-2p::GFP] (vdac-1 oe)</i> 8x	This study	N/A
<i>C. elegans</i> : MGH357 <i>glp-1(e2141);alxEx91[vdac-1p::VDAC-1::FLAG::SL2::GFP]</i>	This study	N/A
<i>C. elegans</i> : MGH167 <i>sid-1(qt9);alxS9[vha-6p::SID-1::SL2::GFP]</i>	This study	N/A
<i>C. elegans</i> : MGH206 <i>riect-1(mg451);sid-1(qt9);alxS9 [vha-6p::SID-1::SL2::GFP]</i>	This study	N/A
<i>C. elegans</i> : MGH369 <i>sgk-1(mg455); h1h-30p::HLH-30::GFP</i>	This study	N/A
<i>C. elegans</i> : MGH371 <i>riect-1(mg451); adls2122 [lgg-1p::GFP::lgg-1 + rol-6(su1006)]</i>	This study	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<i>C. elegans</i> : MGH372 <i>sgk-1(mg455);adls2122 [lgg-1p::GFP::lgg-1 + rol-6(su1006)]</i>	This study	N/A
<i>C. elegans</i> : MGH392 <i>mgIs60[sgk-1::GFP pRF4]; alxIs28[vdac-1p::VDAC-1::FLAG; myo-2p::GFP]</i>	This study	N/A
<i>C. elegans</i> : MAH885 <i>sgk-1(mg455);unc-119(ed3)? III; ppls?[unc-119(+); lgg-1p::gfp::lgg-1(G116A)]</i>	This study	N/A
<i>C. elegans</i> : MAH886 <i>sgk-1(mg455);unc-119(ed3)? III; ppls?[unc-119(+); lgg-1p::gfp::lgg-1(G116A)]</i>	This study	N/A
Mouse: B6.Cg-Sgk1[flox/flox]	Fejes-Tóth et al., 2008	N/A; RRID: MGI:5317851
Mouse: B6.Cg-Speer6-ps1 ^{Tg(Alb-cre)21Mgn/J}	The Jackson Laboratory	003574; RRID: IMSR_JAX: 003574
Mouse: B6.Cg-SgkTKO	This study	N/A
Oligonucleotides		
<i>Sgk1</i> sgRNA targeting sequence: TAAGCAGCCGTAT GACCGGA	This paper	N/A
<i>Sgk2</i> sgRNA targeting sequence: CGGAGCCTCTA CGCCGTGA	This paper	N/A
<i>Sgk3</i> sgRNA targeting sequence: CAAGGCACTG GCGATCTCCG	This paper	N/A
QPCR, <i>act-3</i> , Forward: TGCGACATTGATATCCG TAAGG	This paper	N/A
QPCR, <i>act-3</i> , Reverse: GGTGGTTCCTCCGAAAGAA	This paper	N/A
QPCR, <i>nd-1</i> , Forward: AGCGTCATTTATTGGGAA GAAGAC	This paper	N/A
QPCR, <i>nd-1</i> , Reverse: AAGCTTGTGCTAATCCCAT AAATGT	This paper	N/A
QPCR, <i>IL-10</i> , Forward: GCTCTTACTGACTGGCA TGAG	This paper	N/A
QPCR, <i>IL-10</i> , Reverse: CGCAGCTCTAGGAGCATGTG	This paper	N/A
QPCR, <i>IL-1β</i> , Forward: AAATACCTGTGGCCTTGGGC	This paper	N/A
QPCR, <i>IL-1β</i> , Reverse: CTTGGGATCCACACTC TCCAG	This paper	N/A
QPCR, <i>bec-1</i> , Forward: ACGAGCTTCATTCGC TGGAA	This paper	N/A
QPCR, <i>bec-1</i> , Reverse: TTCGTGATGTTGTACGCCGA	This paper	N/A
QPCR, <i>lgg-1</i> , Forward: GCCGAAGGAGACAAG ATCCG	This paper	N/A
QPCR, <i>lgg-1</i> , Reverse: GGTCTGGTAGAGTTGTCCC	This paper	N/A
QPCR, <i>unc-51</i> , Forward: CGCCGGTGGTTCAGC GGATT	This paper	N/A
QPCR, <i>unc-51</i> , Reverse: TATCCTGGGTGTCGGC GGGG	This paper	N/A
QPCR, <i>h1h-30</i> , Forward: GGATTAAGGAGCTCGG ACAA	This paper	N/A
QPCR, <i>h1h-30</i> , Reverse: TTTTGGAGCACTCGGATGT	This paper	N/A
Recombinant DNA		
GST-SGK-1	This paper	N/A
VDAC-1-Myc-His	This paper	N/A
Flag-SGK1-WT	This paper	N/A
pCMV5-Flag	This paper	N/A
Flag-SGK1-S422D	This paper	N/A
Flag-SGK1-S422A	This paper	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mus-VDAC1-Myc	This paper	N/A
Ce-VDAC-1	This paper	N/A
Mus-VDAC1-Myc S13A	This paper	N/A
Mus-VDAC1-Myc S35A	This paper	N/A
Mus-VDAC1-Myc T107V	This paper	N/A
Mus-VDAC1-Myc T116V	This paper	N/A
Mus-VDAC1-Myc S137A	This paper	N/A
Mus-VDAC1-Myc S193A	This paper	N/A
Mus-Flag-VDAC1	This paper	N/A
Mus-Flag-VDAC1 S101/102A	This paper	N/A
Mus-Flag-VDAC1 S104A	This paper	N/A
HA-Ub	Johnathan R. Whetstone's lab	N/A
Software and Algorithms		
ImageJ2	NIH	https://imagej.nih.gov/ij/
GraphPad Prism7	GraphPad Software	https://www.graphpad.com/
Oasis2	Han et al., 2016 Oncotarget 11269	https://sbi.postech.ac.kr/oasis2/history/

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Alexander A. Soukas (asoukas@mgh.harvard.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Nematode Strains

N2 Bristol (RRID:WB-STRAIN:N2_(ancestral)) was used as the wild-type strain. The listed mutant strains were used: MGH12 *mgIs60* [*sgk-1::GFP* pRF4], MGH48 *mgIs43*[*ges-1p::GFP*_PTS1], MGH266 *rict-1(mg451)* 8x, MGH300 *sgk-1(mg455)* 8x, MGH55 *daf-16(mgDf47)* 3x, MGH361 *atfs-1(tm4525)* 3x, TK22 *mev-1(kn1)* 5x (RRID:WB-STRAIN:TK22), DA465 *eat-2(ad465)* (RRID:WB-STRAIN:DA465), CB4037 *glp-1(e2141)* (RRID:WB-STRAIN:CB4037), MGH358 *mev-1(kn1);sgk-1(mg455)*, MGH359 *atfs-1(tm4525);sgk-1(mg455)*, MGH360 *daf-16(mgDf47);sgk-1(mg455)*, MGH369 *sgk-1(mg455)*, DA2123 *adIs2122* [*lgg-1p::GFP::lgg-1 + rol-6(su1006)*] (RRID:WB-STRAIN:DA2123), MGH371 *rict-1(mg451); adIs2122* [*lgg-1p::GFP::lgg-1 + rol-6(su1006)*], MGH372 *sgk-1(mg455); adIs2122* [*lgg-1p::GFP::lgg-1 + rol-6(su1006)*], MGH392 *mgIs60*[*sgk-1::GFP* pRF4]; *alxIs28*[*vdac-1p::VDAC-1::FLAG myo-2p::GFP*], RD202 *unc-119(ed3)* III; *ppls??[unc-119(+)* *lgg-1p::gfp::lgg-1(G116A)*], MAH885 *sgk-1(mg455);unc-119(ed3)?* III; *ppls?[unc-119(+)* *lgg-1p::gfp::lgg-1(G116A)*]; JIN1821 *jInIs[hLh-30p::HLH-30::GFP]*, N2, TK22 *mev-1(kn1)*, DA465 *eat-2(ad465)* and CB4037 *glp-1(e2141)* strains were obtained from the *Caenorhabditis* Genetics Center (CGC). MGH361 *atfs-1(tm4525)* was a gift from Cole Haynes' lab. JIN1821 *jInIs[hLh-30p::HLH-30::GFP]* was a gift from Javier Irazoqui. For mitochondrial targeted RFP expression: MGH84 *mgIs54*[*myo-3p::mito::mRFP3*] and MGH353 *sgk-1(mg455);mgIs54*[*myo-3p::mito::mRFP3*] were used. For *vdac-1* overexpression, MGH354 *alxEx91*[*vdac-1p::VDAC-1::FLAG::SL2::GFP*] (*vdac-1 oe1*), MGH355 *alxEx92*[*vdac-1p::VDAC-1::FLAG myo-2p::GFP*] (*vdac-1 oe2*), MGH356 *alxIs28*[*vdac-1p::VDAC-1::FLAG myo-2p::GFP*] (*vdac-1 oe*) 8x and MGH357 *glp-1(e2141);alxEx91*[*vdac-1p::VDAC-1::FLAG::SL2::GFP*] were generated for this study. For tissue-specific RNAi, MGH167 *sid-1(qt9);alxIs9*[*vha-6p::SID-1::SL2::GFP*], MGH206 *rict-1(mg451);sid-1(qt9);alxIs9*[*vha-6p::SID-1::SL2::GFP*] were used. Nematode strains were maintained on nematode growth medium (NGM) plates seeded with *E. coli* OP50. All experiments were conducted at 20°C unless otherwise noted.

E. coli Strains

Non-RNAi experiments were all conducted on NGM plates containing *E. coli* OP50-1 (CGC) as the food source. Cultures of *E. coli* OP50 were grown in LB broth for 15h at 37°C without shaking and seeded directly onto NGM plates. Plates were used 3-7 days after seeding.

All RNAi experiments were conducted using *E. coli* HT115 (DE3) bacteria (Ahringer and Vidal RNAi Library) as the food source. Cultures of *E. coli* HT115 (DE3) were grown in LB broth for 15h at 37°C with shaking, then cultures were concentrated to 1/10 volume by centrifugation and seeded directly onto NGM plates containing 5 mM isopropyl-B-D-thiogalactopyranoside and 200 µg/ml carbenicillin. Plates were used 1-5 days after seeding.

Mice

All mice were housed in a temperature-controlled room under a 12 hr light–dark cycle and under specific pathogen-free conditions, and provided *ad libitum* access to water and standard laboratory chow (Prolab Isopro RMH 3000). *Sgk1[flox/flox]* (RRID: MGI:5317851) mice were as we previously published (Fejes-Tóth et al., 2008). *Sgk1^{Lko}* mice were generated by crossing Albumin-Cre mice (The Jackson Laboratory, RRID: IMSR_JAX: 003574) with *Sgk1[flox/flox]* mice. Double *Sgk2;Sgk3* global knockout mice were generated by C57BL/6 zygotic injection of small guide RNAs targeting *Sgk2* and *Sgk3* and *S.p.Cas9* mRNA at the Genome Modification Facility, Harvard University. Mice were backcrossed four times prior to experimentation. *SgkTKO* mice were generated by crossing *Sgk1^{Lko}* mice to *Sgk2;Sgk3* global knockout mice. Twelve-week-old male *Sgk1^{Lko}* and control mice, and 24-week-old male *SgkTKO* and control mice were used for hepatic ischemia/reperfusion assays. All animal procedures were approved under protocol 2010N000186 by Massachusetts General Hospital Subcommittee on Research Animal Care. Guide RNA sequences used for generating double *Sgk2;Sgk3* global knockout mice are as follows (See also Table S2):

Sgk2: CGGAGCCTTCTACGCCGTGA;

Sgk3: CAAGGCACTGGCGATCTCCG.

Cell Lines, Culture Conditions and Transfection

The AML12 (alpha mouse liver 12) cell line was established from hepatocytes from a male mouse (CD1 strain, line MT42) transgenic for human TGF alpha (ATCC, ATCC® CRL-2254). AML12 hepatocytes were grown in a 1:1 mixture of DMEM and Ham's F12 medium (Thermo Fisher) with 0.005 mg/ml insulin, 0.005 mg/ml transferrin, 5 ng/ml selenium, 40 ng/ml dexamethasone, 10% fetal bovine serum (FBS, Sigma) and 1% penicillin-streptomycin (PS, Life Technologies), in a 5% CO₂ atmosphere at 37°C. Primary cultured mouse hepatocytes were prepared from male C57BL/6, liver-specific *Sgk1* knockout mice (genotype Albumin-Cre;*Sgk1[flox/flox]*) and their wild-type littermate controls (genotype *Sgk1[flox/flox]*) by collagenase perfusion method as described previously (Li et al., 2010), and were grown in Williams' Medium E supplemented with 10% FBS and 1% PS. All animal procedures were approved under protocol 2010N000186 by Massachusetts General Hospital Subcommittee on Research Animal Care. HEK293T cells (a gift from Joseph Avruch's lab), generated from female human embryonic kidney with stably expressing the SV40 T-antigen, were grown in high glucose DMEM (Thermo Fisher) supplemented with 10% FBS and 1% PS in a 5% CO₂ atmosphere at 37°C. For transfection, Lipofectamine 3000 was used according to the instructions of manufacturer (Thermo Fisher).

METHOD DETAILS

Generation of *Sgk1* Knockout AML12 Cells

Sgk1 knockout AML12 cells were generated using CRISPR/Cas9 (Cong et al., 2013) with modifications. The px330 vector expressing Cas9 and guide RNA was electro-transfected into AML12 cells together with a plasmid conferring puromycin resistance. Forty-eight hours after transfection, puromycin was added to the cells at 5 μg/ml for 48 hours, then puromycin resistant cells were trypsinized, counted and diluted to 10 cells/ml, then 100 μL of the diluted cell mixture per well was seeded into a 96 well plate, and single-clones were expanded to generate stable cell lines. *Sgk1* knockout was confirmed by western blotting. Primers used for sgRNA expression are (See also Table S2): sense, 5'-CACCGTAAGCAGCCGATGACCGGA-3', antisense, 5'-AAACTCCGGTCATACGGCTGCTTAC-3'.

Transmission Electron Microscopy (TEM)

C. elegans TEM was performed by the Transmission Electron Microscopy Core at Massachusetts General Hospital (Lapierre et al., 2013). Wild-type and *sgk-1(mg455)* synchronized L4 populations were collected and incubated in fixation buffer (2.5% glutaraldehyde and 1.0% paraformaldehyde in 0.05M sodium cacodylate at pH7.4 plus 3.0% sucrose). During the initiation of fixation, animals were cut in half with a surgical blade in a drop of fixative under a dissecting microscope, fixed overnight at 4°C, rinsed in 0.1M cacodylate, post-fixed in 1.0% osmium tetroxide 0.1M cacodylate buffer, rinsed in buffer and water. Animals were embedded in 2% agarose in phosphate buffered saline (PBS), dehydrated through a graded series of ethanol washes to 100%, then 100% propylene oxide, and finally 1:1 propylene oxide/EPON overnight. Blocks were infiltrated in 100% EPON and then embedded in fresh EPON overnight at 60°C. Thin sections were cut on a Leica UC6 ultramicrotome and collected on formvar-coated grids. Sections were post-stained with uranyl acetate and lead citrate and viewed using a JEOL 1011 transmission electron microscope at 80 kV with an AMT digital imaging system (Advanced Microscopy Techniques). Autolysosomes were identified as vesicles containing material undergoing degradation in micrographs of a cross-section of the intestine.

Measuring Autophagy Levels in *C. elegans* and Cells

Autophagy levels were assessed following published guidelines (Klionsky et al., 2016; Zhang et al., 2015). Autophagosomes were scored as follows (Lapierre et al., 2013). For *C. elegans* starvation, worms were kept in M9 for 24h in a 3cm dish with gentle shaking. For imaging and puncta quantification, animals were mounted on a 2% agarose pad in M9 medium containing 0.1% NaN₃ and GFP::LGG-1/Atg8 puncta were counted using a Zeiss Imager Z1 including apotome.2 with a Hamamatsu orca flash 4LT camera and Zen 2.3 software. The total number of GFP::LGG-1/Atg8 puncta was counted in all visible hypodermal seam cells, the striated

body-wall muscle, the three to four most proximal intestinal cells or the terminal pharyngeal bulb at 100x magnification. For mammalian cells, autophagosomes were stained using Cyto-ID according to manufacturer instructions (Enzo Life Sciences). Cells were fixed with 4% paraformaldehyde for 20 min and rinsed with PBS twice. Cells were mounted and visualized under a Leica DM6000 microscope with a 63X oil immersion objective.

Chloroquine *C. elegans* Treatment

For chloroquine treatment, 100 mM chloroquine was added on top of the NGM plates seeded with OP50. About 500 synchronized L4 animals were seeded onto these NGM plates for 18h before harvest.

Western Blotting

Worm and cell lysates were prepared in RIPA buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton x-100, 1% Sodium deoxycholate, 0.1% SDS, 1 mM EDTA with proteinase inhibitors cocktail from Roche) unless otherwise indicated, followed by water bath sonication on maximum energy in a Diagenode Bioruptor XL 4°C water bath at 30 s/30 s on and off intervals for a total of 15 min. Lysates were cleared of insoluble material by centrifugation at 21,000 g at 4°C and the supernatant was retained for western blotting. Protein concentration was determined using the Pierce BCA assay (Thermo Fisher). SDS-PAGE was conducted followed by electrophoretic transfer to nitrocellulose membrane at 100 V for 1 hour at 4°C. Immunoblots were performed according to primary antibody manufacturers' protocols. For immunodetection of primary antibodies, goat-anti-rabbit-HRP conjugate or goat-anti-mouse-HRP conjugate (GE Healthcare) was used at 1:5,000 in 5% BSA dissolved in TBST, and HRP was detected using West-Pico chemiluminescence substrate (Thermo Pierce). The western blot results shown are representative of at least three independent experiments.

Longevity Assay

Lifespan analysis was conducted at 20°C except where indicated, as previously published (Soukas et al., 2009). Synchronized L1 animals were seeded onto NGM (for mutants and CsA treatment) or RNAi plates and allowed to grow until the L4 stage (for RNAi). On day 0 as indexed in the figure legend, about 100 L4 worms per plate were transferred onto NGM or RNAi plates. All plates were supplemented with 100 μM 5-fluorodeoxyuridine (FUDR) to suppress progeny production. For CsA treatment, about 100 synchronized L1 animals were seeded onto NGM plates containing 5 μM CsA, and at the L4 stage, worms were transferred to NGM plates containing both CsA and FUDR for the remainder of their life. For *glp-1* mutant worms, synchronized L1 animals were raised at 25°C (to eliminate germ cells) until young adulthood, then worms were shifted to 20°C for the remainder of life. When compared with other strains using FUDR, strains carrying the *glp-1* mutation were also put on plates with FUDR for lifespan analysis. Dead worms were counted every day. Statistical analysis was performed with online OASIS2 resources (Han et al., 2016).

RNA Interference (RNAi)

RNAi clones were isolated from a genome-wide *E. coli* RNAi library, sequence verified, and fed to animals. RNAi feeding plates (6 cm) were prepared using a standard NGM recipe with 5 mM isopropyl-B-D-thiogalactopyranoside and 200 μg/ml carbenicillin. RNAi clones were grown for 15 hours in Luria Broth (LB) containing 200 μg/ml carbenicillin with shaking at 37°C. The stationary phase culture was then collected, concentrated through centrifugation, the supernatant was discarded and the pellet was resuspended in LB to 10% of the original culture volume; 300 μL of each RNAi clone concentrate was added to RNAi plates and allowed to dry no more than 48 hours prior to adding the worm embryos or animals.

Quantitative RT-PCR

Worms and mouse tissues were flash frozen in liquid nitrogen and kept in –80°C until RNA preparation. To quantify changes in mRNA abundance in nematode and mammalian cells, total RNA was extracted using RNAzol RT (Molecular Research Center) according to manufacturer instructions. RNA was treated with RNase free DNase prior to reverse transcription with the Quantitect reverse transcription kit (QIAGEN). Quantitative RT-PCR was conducted in triplicate using a Quantitect SYBR Green PCR reagent (QIAGEN) following manufacturer instructions on a Bio-Rad CFX96 Real-Time PCR system (Bio-Rad). For *C. elegans*, all assays were performed with 1000 animals per sample at late L4 stage or as indicated in the figure legend. For mouse tissue samples, liver samples were homogenized in RNAzol using a TissueLyser II (QIAGEN), centrifuged for 10 min at 12,000 g to pellet debris, and the supernatant was collected and subjected to RNA extraction per the manufacturer protocol. Expression levels of tested genes were presented as normalized fold changes to the mRNA abundance of control genes indicated in the figures by the $\delta\delta$ Ct method.

The primers used for qPCR are as follows (See also Table S2):

act-3, Forward: TGCGACATTGATATCCGTAAGG
act-3, Reverse: GGTGGTTCCTCCGAAAGAA
nd-1, Forward: AGCGTCATTTATTGGGAAGAAGAC
nd-1, Reverse: AAGCTTGTGCTAATCCATAAATGT
IL-10, Forward: GCTCTTACTGACTGGCATGAG
IL-10, Reverse: CGCAGCTCTAGGAGCATGTG
IL-1β, Forward: AAATACCTGTGGCCTTGGGC

IL-1 β , Reverse: CTTGGGATCCACACTCTCCAG
bec-1, Forward: ACGAGCTTCATTCGCTGGAA
bec-1, Reverse: TTCGTGATGTTGTACGCCGA
lgg-1, Forward: GCCGAAGGAGACAAGATCCG
lgg-1, Reverse: GGTCCTGGTAGAGTTGTCCC
unc-51, Forward: CGCCGGTGGTTCAGCGGATT
unc-51, Reverse: TATCCTGGGTGTCGGCGGGG
hlh-30, Forward: GGATTAAGGAGCTCGGACAA
hlh-30, Reverse: TTTTGGAGCACTCGGATGT

Brood Size

Ten synchronized animals per strain were transferred daily to fresh RNAi plates containing *E. coli HT115* expressing target dsRNAs, and hatched progeny were counted 3 days later. Total brood size was determined by adding progeny produced across all days.

Developmental Timing

Developmental timing analysis was conducted on synchronized L1 animals. Synchronous animals were dropped onto RNAi plates containing empty vector RNAi control or a sequence-verified target RNAi in *E. coli HT115* and grown at 20°C. Hermaphrodite animals were scored for their transition into adulthood by appearance of the vulvar slit at the indicated time points. Two to three biological replicates were carried out for each condition.

Nile-Red Staining

Fixative Nile-Red staining was conducted by washing 200–300 young adult animals from synchronized L1 worms obtained by egg prep. The worms were resuspended and washed twice with PBS and then suspended in 40% isopropanol for 5 min. Then the worms were stained in Nile-Red staining solution (3 μ g/ml Nile-Red in 40% isopropanol) in the dark for at least 2 hours. After staining, worms were washed and put in PBS with 0.01% Triton in the dark for 30 min. Imaging and quantitation was then conducted in fluorescence with a GFP/FITC filter using a Leica DM6000 microscope and Leica MM AF software.

SGK-1-GFP Co-IP for Mass Spectrometry

Transgenic *C. elegans* strains that express SGK-1::GFP (MGH12) or GFP only (MGH48) were used for Co-IP. Ten 15cm dishes of mixed age worms were used to get enough protein for mass spectrometry. Briefly, worm lysates were made using RIPA buffer (10mM Tris-HCl pH 7.2, 5mM EDTA, 150mM NaCl, 0.1% SDS, 1.0% Triton X-100, 1% Deoxycholate with Protease inhibitors). SGK-1::GFP was pulled down using an anti-GFP antibody (ThermoFisher) and eluted with 2X Laemmli Buffer without DTT. Samples were loaded onto a 10% SDS-PAGE gel and stained using Coomassie Brilliant Blue. Bands corresponding to broad ranges of molecular weight were cut and sent to The Taplin Biological Mass Spectrometry Facility at Harvard for in gel trypsin digestion and subsequent mass spectrometric peptide/protein identification.

Oxygen Consumption

Synchronized, young adult worms grown on either NGM OP50 or RNAi plates were washed repeatedly with M9 media to remove bacteria. Oxygen consumption of 2500 worms was then measured immediately in 1.5 mL of M9 using a Strathkelvin 928 6-channel Clark electrode respirometer with stirring. Independent measurements were conducted and averaged per strain, per RNAi and normalized to total protein content determined using a BCA assay (Pierce).

ATP Measurement

ATP measurement was carried out according to a modified protocol based on the published literature (Palikaras and Tavernarakis, 2016). 100 synchronized young adult worms were collected in M9 buffer and washed three times. Worm pellets were resuspended with 50 μ L M9 and boiled for 15 min to release ATP and destroy ATPase activity and then spun at 4° at 12,000 \times g for 10 min, and the supernatant was diluted 10 times for ATP measurement. ATP content was measured with an ATP detection kit according to the manufacturer's instructions (ThermoFisher, A22066).

Creation of the VDAC-1::FLAG *C. elegans* Transgenic Lines

For VDAC-1 expression, the entire genomic sequence of the *vdac-1* locus (1367 bp), including introns and exons, plus 2418 bp of promoter were amplified and cloned into a modified fire vector driving VDAC-1 with a C-terminal Flag epitope tag and GFP expression from the same promoter separated by the *gpd-2/3* splice leader sequence (pAS21). The following cloning primers were used: forward (See also Table S2): 5'- ATCGTCGACTCCAGTTGTGGCTTAGAGACTTC -3' and reverse:

5'- CCCGGGCTACTTGTGTCATCGTCTTTGTAGTCGTTGGATGGATCGAATTCGAGTC -3'. The VDAC-1 expression construct was injected at 100 ng/ml into the gonad of wild-type adult animals.

VDAC1 In Vitro Pull-Down Assay and Ubiquitination Assay

C. elegans vdac-1 cDNA was amplified using following primers (See also Table S2):

vdac-1 forward, 5'- CATGCGGCCGCATGGCCCCACCAACCTTCGCTG -3',
vdac-1 reverse, 5'- CGAGATATCGTTGGATGGATCGAATTCGAGT -3',

Amplified fragments were inserted into pcDNA3.1-Myc-His vectors with NotI and EcoRV double digestion. *C. elegans sgk-1* was cloned into the pEBG vector to express a GST-SGK-1 fusion protein using these primers:

forward, 5'- CGCACTAGTGTGAGGAAAGATGAGGTGACAT -3',
reverse, 5'- ATAGCGGCCGCTCAGACCAAACGCGATTGGTG -3'.

Mammalian *Sgk1* and *Vdac1* were cloned into pCMV5-Flag and pcDNA3.1-Myc-His vectors respectively using the following primers:

Sgk1 forward, 5'- ATTTGCGGCCGCTGGAAGATGGTAAACAAAGACA -3',
Sgk1 reverse, 5'- TAGGATCCTCAGAGGAAGGAATCCACAGGA -3',
Vdac1 forward, 5'- CATGCGGCCGCATGGCCGTGCCTCCCACATACG -3',
Vdac1 reverse, 5'- CGAGATATCTGCTTGAATTCCAGTCCTAGG -3'.

To make the constitutively active SGK1, Ser422 was changed to Asp by site directed mutagenesis, a change previously shown to increase kinase activity (Park et al., 1999).

For GST-pulldown assays, GST-SGK-1 protein was purified from HEK293T cells transfected with expression vectors using the Pierce GST Spin Purification Kit (Thermo Scientific, Prod# 16106). Five micrograms of purified GST and GST-SGK-1 proteins were re-immobilized with pre-equilibrated 20 μ L GST beads in lysis buffer. VDAC-1-Myc-His protein was purified from HEK293T cells following transfection of the indicated expression vectors using the Pierce HisPur Ni-NTA Purification Kit (Thermo Scientific, Prod# 88227). Protein concentration was determined with the Pierce BCA Protein Assay Kit (Thermo Scientific) before the GST-pulldown assay. Five micrograms of His purified protein were incubated with the immobilized beads at 4°C for 1.5 hours. After washing the beads three times with lysis buffer, samples were analyzed by western blot using the anti-Myc antibody.

For the Co-IP assay with overexpressed protein, GST-SGK-1 and VDAC-1-Myc-His were co-transfected into HEK293T cells. Forty-eight hours after transfection, cells were lysed using Co-IP buffer (0.025 M Tris, 0.15 M NaCl, 0.001 M EDTA, 1% NP-40, 5% glycerol; pH 7.4), GST-SGK-1 protein was pulled down using the Pierce GST Spin Purification Kit, and samples were analyzed by western blot using anti-Myc antibody. Similarly, for mammalian SGK1 and VDAC1, Flag-SGK1 and VDAC1-Myc-His were co-transfected into HEK293T cells for 48h, cells were then lysed using Co-IP buffer, and lysate was incubated with M2-Flag magnetic beads from SIGMA following its Co-IP protocol. Western blot was used for detection of VDAC1-Myc. For the Co-IP assay of mammalian endogenous protein, 10⁷ HEK293T cells were lysed using Co-IP buffer, SGK1 or VDAC1 protein was pulled down using SGK1 antibody (Cell signaling Technology) and protein A magnetic beads (New England Biolabs) or VDAC1 antibody (Santa Cruz) and protein G magnetic beads (New England Biolabs) following the manufacturer's protocol.

For detection of VDAC1 ubiquitination, Flag-SGK1, VDAC1-Myc-His and HA-Ub plasmids were co-transfected into HEK293T cells for 48h, then cell lysates were made using RIPA buffer, and magnetic Protein A beads (New England Biolabs) were used for Myc IP following the manufacturer's protocol. Anti-HA antibody was used to detect the ubiquitination of VDAC1 protein.

mtDNA Copy Number

The mtDNA copy number was measured by quantitative PCR. Worms at the indicated developmental stage were singled in worm lysis buffer for PCR. Primers for NADH dehydrogenase subunit 1 (nd1) and actin-3 (act-3) were used in determination of mtDNA copy number. Primers used were as follows (See also Table S2): act-3 forward primer: 5'- TGCGACATTGATATCCGTAAGG -3', and reverse primer 5'- GGTGGTTCCTCCGAAAGAA -3'; nd1 forward primer 5'- AGCGTCATTTATTGGGAAGAAGAC -3', and reverse primer 5'- AAGCTTGTGCTAATCCATAAATGT -3'. PCR conditions were 2 min at 50°C, 10 min at 95°C, followed by 40 cycles of 15 s at 95°C and 60 s at 60°C. Quantitative PCR was conducted in triplicate using Quantitect SYBR Green PCR reagent (QIAGEN) following manufacturer instructions on a CFX96 Real-Time PCR system (Bio-Rad). The C_T values of nd1 were normalized to nuclear genomic DNA (act-3). Quantitative PCR was performed for each initial template copy number at least four times and the results were reproducible.

Mitochondrial Ca²⁺ Uptake

Cells were permeabilized in a buffer containing 125 mM KCl, 2 mM K₂HPO₄, 1 mM MgCl₂, 20 mM HEPES pH 7.2, 0.005% digitonin, 5 mM glutamate and malate, and a cell-impermeable Ca²⁺ indicator (0.5 μ M Calcium Green-5N). The fluorescence of the Ca²⁺ indicator dye was monitored in real time using a PerkinElmer LS55 fluorimeter using a cuvette with stirring at room temperature. Pulses of CaCl₂ were given sequentially (5 μ l of 3 mM CaCl₂ was added to 500 μ l assay buffer). At least three replicates were performed for each experiment, and representative traces are shown.

Paraquat Treatment

For paraquat treatment, 8 mM paraquat was added on top of the NGM (for mutants) or RNAi agar plates. About 100 synchronized L4 animals were seeded onto the NGM or RNAi plates seeded with OP50 or HT115 expressing dsRNA for RNAi respectively, and dead worms were counted every day. Statistical analysis was performed with online OASIS2 resources.

Identification of the VDAC1 Phosphorylation Site by Mass Spectrometry

Proteins were separated by SDS-PAGE, gel bands were cut and destained in 50 mM ammonium bicarbonate:acetonitrile (50:50) until clear. Protein were in-gel reduced with dithiothreitol (DTT), alkylated with iodoacetamide, and digested with trypsin. Peptides were extracted and subjected to C18 solid phase extraction. The generated peptides were subjected to microcapillary liquid chromatography followed by tandem mass spectrometry (LC-MS2) on an Orbitrap Fusion Lumos mass spectrometer (Thermo Scientific). Full MS spectra were acquired using the Orbitrap analyzer, and MS2 spectra were acquired in a data dependent mode after CID fragmentation using the linear ion trap. MS2 spectra were assigned using a SEQUEST proteomics analysis platform requiring cysteine residues to be carbamidomethylated (mass increment of 57.02146 Da), while methionine residues were allowed to be oxidized (15.99492 Da) and serine, threonine and tyrosine residues were allowed to be phosphorylated (79.96633 Da). The database used in the search was the UniProt database (02/04/2014) of human protein sequences including known contaminants such as trypsin. Based on the target-decoy database search strategy and employing linear discriminant analysis and posterior error histogram sorting, peptide and protein assignments were filtered to false discovery rate (FDR) of < 1%. The localization of phosphorylation sites was determined using the modscore algorithm (Schwartz and Gygi, 2005). For quantification, the MS1 signal for the phosphorylated peptide and unphosphorylated peptide was integrated using Qual Browser software (Buisson et al., 2017). Integration was done for VDAC1 protein incubated with or without SGK1 S422D in duplicate and statistical significance confirmed by Student's t test.

Hepatic Ischemia/Reperfusion Model

The hepatic warm I/R model was performed as described below. Under ketamine (100 mg/kg body weight, administered intraperitoneally [IP]) and xylazine (10 mg/kg IP) anesthesia, the liver was exposed, and the left and median lobes were clamped with a microvascular clamp for 45 minutes. Ischemia was confirmed by visualizing the blanching of the ischemic lobes. The clamp was removed, and reperfusion confirmed on the basis of immediate color change before the abdomen was closed with continuous silk suture. Absence of ischemic color change or lack of reperfusion resulted in exclusion from further analysis. Temperature was maintained at 37°C by a warming pad. Six hours after reperfusion, mice were sacrificed by exsanguination after being anesthetized with ketamine and xylazine. For Cyclosporine A pretreatment, a dose of 10 mg/kg CsA were injected IP twice at 12h and 1h before ischemia.

QUANTIFICATION AND STATISTICAL ANALYSIS

All western blotting quantifications were conducted in ImageJ. Statistical analyses were performed using Prism (GraphPad Software). The statistical differences between control and experimental groups were determined by two-tailed Student's t test (two groups), one-way ANOVA (more than two groups), or two-way ANOVA (two independent experimental variables), with corrected *P* values < 0.05 considered significant. Asterisks denote corresponding statistical significance **p* < 0.05; ***p* < 0.01; ****p* < 0.001 and *****p* < 0.0001. The statistical tests performed and definition of "n" numbers in this study are indicated in the figure legends. For Figures 1A, 1B, 2H, S1C, S2B–S2F, and S3K, "n" means the number of worms. For Figures 7B–7H, "n" means the number of mice. For Figures 1C, 1F, 1G, 2E, and 5B, "n" means the number of cells that were analyzed. For Figures 1D, 1H, 1I, 1J, 3J, 3K, 4A–4C, 4G–4K, 5A, 5C–5G, 5I, 6A–6C, S1A, S1B, S1E, S2K, S3G, S3H, S3J, S6D–S6F, S6L, S7B, and S7C, "n" means biological replicates. For *C. elegans* western blotting, oxygen consumption, and ATP determination, each sample within each biological replicate corresponds to a sample pooled from 100–1000 animals. For cell culture western blotting, each sample within each biological replicate corresponds to one well from a tissue culture plate. For primary hepatocyte experiments, each biological replicate corresponds to hepatocytes isolated from one mouse per genotype, divided evenly between experimental conditions. For mouse liver western blotting (Figures 7D and 7H) each sample corresponds to protein extract from one mouse. The log rank test was used to determine significance in lifespan analyses using online OASIS2. For details of survival analyses and statistics, see Table S1.




Figure S1. SGK1 Regulates Autophagy in Both *C. elegans* and Mouse Primary Hepatocytes, Related to Figure 1

(A) SGK-1 protein level is decreased under starvation. SGK-1::GFP fusion protein is detected by western blotting for GFP. (n = 3, Student's t test)

(B) Phosphorylation of *C. elegans* α AMPK(Thr172) in *sgk-1* mutant and WT animals, as assessed by western blot. (n = 3, Student's t test)

(C) Brood size of individual *sgk-1* mutant and WT *C. elegans* as determined daily from day 1 of adulthood for five days. (n = 10 per genotype, two-way ANOVA)

(D) Imaging *in situ* using the fluorescent autophagosome indicator Cyto-ID indicates elevated autophagy levels in AML12 *Sgk1*^(-/-) cells. Scale bar: 20 μ m.

(E) Immuno-blotting of LC3A/B turnover in control (littermates without the *Cre* transgene, *Sgk1*^{fl^{ox}/fl^{ox}}) and *Sgk1*^{Lko} primary hepatocytes under full medium conditions. (n = 4 replicates from 2 mice per group, Student's t test)

(F) The fluorescent autophagosome indicator Cyto-ID demonstrates elevated autophagy levels in *Sgk1*^{Lko} primary hepatocytes both under full medium and HBSS starvation conditions, scale bar: 20 μ m.

*p < 0.05, **p < 0.01. All bars indicate means and SEM.




Figure S2. SGK-1 Modulates Lifespan, Development, and Reproduction through Altered Levels of Autophagy, Related to Figure 2

(A) RNAi of *bec-1* restores normal lifespan in both *sgk-1* and *rict-1;sgk-1* double mutants (log rank test).

(B) RNAi of *unc-51* increases brood size in *sgk-1* mutants ($n = 5$ per group, Student's *t* test).

(C and D) RNAi of *bec-1*, *lgg-1*, and *unc-51* accelerates growth rate in *sgk-1* mutants. Developmental stage was evaluated 61 hours after seeding of synchronized *sgk-1* L1 worms (C) or 48 hours after seeding of synchronized WT L1 worms (D). YA: young adult; L4: larval stage 4 ($n > 50$ per genotype, per replicate for 3 replicates, two-way ANOVA).

(E) RNAi of *bec-1*, *lgg-1*, and *unc-51* does not significantly change fat mass in *sgk-1* mutants. Fixative Nile red staining was used to assess fat levels. ($n = 69, 108, 70$ and 61 animals for ev *bec-1*, *lgg-1*, and *unc-51* RNAi, respectively).

(F) Intestine specific RNAi of *bec-1*, *lgg-1*, and *unc-51* accelerates growth rate in *rict-1* mutants. Developmental stage was checked 62 hours after seeding of synchronized intestine specific RNAi *rict-1* L1 worms ($n > 60$ per genotype, per replicate for 3 replicates, two-way ANOVA).

(G and H) Intestine-specific RNAi of *bec-1* partially rescues shortened lifespan of *rict-1* mutant worms (G) but slightly shortens lifespan of WT worms (H) (log rank test).

(I) RNAi of *bec-1* beginning at the L1 stage of larval development partially restores normal lifespan in *sgk-1* mutants compared to whole life (wl) RNAi (log rank test).

(J) RNAi of *bec-1* beginning in adulthood is insufficient to restore normal lifespan in *sgk-1* mutants (log rank test).

(K) Comparison of *bec-1* RNAi efficiency in *sgk-1* mutant worms only during larval stages (larval) and started from young adulthood for three days (adult) ($n = 4$, two-way ANOVA).

For tabular survival data and biological replicates see also Table S1. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. All bars indicate means and SEM.




Figure S3. SGK-1-Dependent Regulation of mPTP Function Impacts Lifespan, Related to Figure 3

(A) Upon co-transfection of mouse Flag-SGK1 and VDAC1-Myc-His in 293T cells, VDAC1-Myc-His co-immunoprecipitates with Flag-SGK1.
 (B) Interaction between VDAC1 and SGK1 is not regulated by wortmannin treatment.
 (C) The mPTP opens earlier in mouse *Sgk1*($-/-$) AML12 cells versus WT cells under a calcium challenge. Two million cells were used for this experiment, versus ten million in Figure 3D.
 (D) RNAi of *atp-5* extends lifespan in *sgk-1* mutants (log rank test).
 (E) mPTP inhibitor CsA treatment partially restores normal lifespan in *sgk-1* mutants. No obvious effects are seen in WT animals (log rank test).
 (F) Intestine specific RNAi of *ant-1.1*, similar to *bec-1*, partially restores normal lifespan in short-lived *rict-1* mutant worms (log rank test).
 (G) ATP level is decreased in both *rict-1* and *sgk-1* mutants compared with wt animals ($n = 4$, one-way ANOVA).
 (H) Mitochondrial DNA copies in WT N2, *rict-1* and *sgk-1* mutants as measured by qPCR. *nd-1* was used as mitochondrial DNA marker normalized to nuclear DNA *act-3* ($n = 7$).
 (I) Representative image of mitochondrially localized RFP in muscle of WT N2 and *sgk-1* mutants, scale bar: 200 μm .
 (J) Increased SGK-1 protein levels under mitochondrial stress induced by RNAi to electron transport chain genes, as indicated by western blotting of SGK-1::GFP. Cell lysates were made using SDS lysis buffer ($n = 3$, one-way ANOVA).
 (K) Representative image of the SGK-1::GFP reporter under mitochondrial stress ($n = 45, 23, 29$ and 25 animals for *ev*, *nuo-1*, *nuo-6* and *cco-4* RNAi, respectively, one-way ANOVA), scale bar: 200 μm .
 For tabular survival data and biological replicates see also Table S1. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. All bars indicate means and SEM.




Figure S4. SGK1 Regulates VDAC1 Protein Level, Related to Figure 4

(A) VDAC-1::Flag fusion protein detected by western blot using Flag antibody in *C. elegans* overexpressing VDAC-1 (*vdac-1 oe*).

(B) Extrachromosomal expression of VDAC-1::Flag confirmed by western blotting in the two *vdac-1* overexpression lines.

(C) ANT protein level measured by western blot in WT, *rict-1* and *sgk-1* mutant *C. elegans*.

(D) ANT1 and VDAC2 protein measured by western blot in WT and *Sgk1*^{Lko} primary hepatocytes.

(E) western blotting of myc-tagged VDAC1 protein with the indicated mutations of candidate Ser and Thr phosphorylation sites with constitutively active SGK1 (S422D) and inactive SGK1 (S422A) co-overexpression.




Figure S5. Increased Mitochondrial Permeability Is Detrimental to Organismal Health, Related to Figure 5

(A) Double RNAi of *bec-1* and *ant-1.1* does not have an additive effect on lifespan extension in *sgk-1* worms (log rank test).

(B and C) Shortened lifespan is also observed in two additional extrachromosomal *vdac-1* overexpression lines compared with WT *C. elegans* (log rank test).

(D) Overexpression of *sgk-1* restores normal lifespan of *vdac-1* overexpression *C. elegans* (log rank test).

(E and F) *rict-1*, *sgk-1* mutants (E) and *vdac-1* overexpression *C. elegans* (F) are more sensitive to paraquat (8mM) treatment compared with WT control (log rank test).

(G-J) Increased resistance to paraquat (8mM) is seen following RNAi of *bec-1* (G and I) or *ant-1.1* (H and J) in wt animals, *sgk-1* and *rict-1* mutants (log rank test).

(K) RNAi of *bec-1* increases paraquat (8mM) resistance in *vdac-1* overexpression *C. elegans* (log rank test).

For tabular survival data and biological replicates see also Table S1. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.




Figure S6. Contribution of Mitochondrial Stress Responses and Mitophagy to *sgk-1* Regulated Changes in Lifespan, Related to Figure 6

(A and B) RNAi of *bec-1* can still restore normal lifespan in *atfs-1(tm4525);sgk-1* (A) and *daf-16(mgDf47);sgk-1* (B) double mutants (log rank test).

(C) RNAi of *skn-1* does not affect the lifespan extension by *bec-1* RNAi in *sgk-1* mutants (log rank test).

(D) *bec-1* RNAi decreases basal and maximal (FCCP) oxygen consumption in WT *C. elegans*. (n = 12, two-way ANOVA).

(E and F) RNAi of *vdac-1* increases basal oxygen consumption in *sgk-1* mutants (E) but has no significant effect on WT *C. elegans* (F) (n = 4, Student's t test).

(G and H) RNAi of mitophagy related genes *pink-1* (G) or *pdr-1* (H) only very modestly extends lifespan of *sgk-1* mutants (log rank test).

(I and J) RNAi of mitochondrial complex II gene *sdhb-1* eliminates autophagy-dependent lifespan extension in both *eat-2(ad465)* mutants (I) and *glp-1(e2141)* mutants (J). For *glp-1(e2141)* lifespan, experiments were conducted at 25 °C (log rank test).

(K) RNAi of *daf-2* still extends lifespan of *vdac-1* overexpression worms (log rank test).

(L) RNAi of *daf-2* decreases VDAC-1 protein level. (n = 3, Student's t test)

For tabular survival data and biological replicates see also Table S1. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. All bars indicate means and SEM.




Figure S7. Increased Sensitivity to Calcium-Induced mPTP Opening in SgkTKO Primary Hepatocytes, Related to Figure 7

(A) The mPTP opens earlier in mouse SgkTKO primary hepatocytes versus control cells when challenged with multiple calcium pulses (arrows). Equal numbers (2×10^6) of cells were used and equivalent biomass was confirmed by western blotting for Actin (right panel).

(B) Increased VDAC1 protein level and LC3 II/LC3 I conversion in SgkTKO primary hepatocytes. (n = 4 replicates for VDAC1 protein level and n = 6 for LC3 II/LC3 I conversion analysis from 3 mice per group, Student's t test).

(C) Decreased phosphorylation levels of p70S6K Thr389 and S6 Ser240/244 in SgkTKO primary hepatocytes. (n = 3, two-way ANOVA).

*p < 0.05, **p < 0.01. All bars indicate means and SEM.